



ALIEN MARINE SPECIES IN NORWAY

Mapping, monitoring and assessment of vectors for introductions



Title (English and Norwegian):

Alien marine species in Norway
Fremmede marine arter i Norge

Subtitle (English and Norwegian):

Mapping, monitoring and assessment of vectors for introductions
Kartlegging, overvåkning og vurdering av vektorer for introduksjoner

Report series:

Rapport fra havforskningen
ISSN:1893-4536

Year - No.:

2022-8

Date:

28.04.2022

Author(s):

Vivian Husa, Ann-Lisbeth Agnalt (IMR), Henrik Berntsen (NINA), Tone Falkenhaus (IMR), Frode Fossøy NINA, Elisabet Forsgren (NINA), Ellen Sofie Grefsrud, Ann Merete Hjelset (IMR), Frank Hanssen (NINA), Even Husby (Bergen havn), Anders Jelmert, Stein Mortensen, Siri Aaserud Olsen (IMR) and Hanno Sandvik (NINA)

Research group leader(s): Bjørn Olav Kvamme (Smittespredning og sykdom) og Kjell Gundersen (Plankton)

Approved by: Research Director(s): Geir Huse Program leader(s): Jan Atle Knutsen

Distribution:

Open

Project No.:

15635-03

On request by:

Miljødirektoratet

Program:

Kystøkosystemer

Research group(s):

Bentiske ressurser og prosesser

Smittespredning og sykdom

Bunnsamfunn

Plankton

Number of pages:

117

Partners

Summary (English):

Norway has the second longest coastline in the world, and it is challenging to monitor non-indigenous marine species (NIMS) along the entire shore including the Norwegian areas in the Barents Sea and along Svalbard. There is currently no national program for such monitoring, however some activity is taking place on specific species and organism groups which is presented here.

Historically transport of NIMS in ballast water have been the main pathway into the Norwegian coast, but with the implementation of the Ballast Water Convention this risk is minimized. Biofouling on vessels coming into the Norwegian coast is thus considered to be the most important vector for marine introduction of new species. An analysis of the frequency and origin (last port call) for 158 000 vessel arrivals into Norwegian ports in the period 2020-2021 is presented. The results show that the Oslofjord area and the west coast is the areas with highest risk for marine introductions by vessels. Other vectors for such introductions into Norway are evaluated like the increasing amount of floating debris which can carry fouling organisms, larvae and eggs to new areas. An analysis of historical data for the established NIMS in Norway show that the southern area of Norway is most susceptible to new species. This pattern is not only dependent on the vector pressure but also reflects the temperature gradient northwards along the coast.

Measures for prevention of new species to arrive and management of problematic species is also discussed.

Summary (Norwegian):

Norge har den nest lengste kystlinjen i verden og det er utfordrende å overvåke fremmede marine arter langs hele kysten og i norske områder i Barentshavet og rundt Svalbard. Det finnes i dag ikke noe nasjonalt program for overvåkning og kartlegging av fremmede marine arter, men det foregår en del aktivitet knyttet til spesifikke arter og organismegrupper som blir presentert her.

Historisk har transport av organismer i ballastvann utgjort den største risikoen for introduksjon av nye marine arter, men med implementeringen av ballastvannsforskriften er denne risikoen blitt betydelig lavere. Påvekst av fremmede organismer på skroget av fartøy som kommer inn til norskekysten fra utenlandske havner trer derfor frem som en av dagens viktigste vektorer for nye introduksjoner. En analyse av frekvensen og opprinnelsen (siste havneanløp) for 158 000 fartøyanløp i norske havner i 2020-2021 presenteres her. Resultatene viser at Oslofjorden og Vestlandet er de områdene med høyest risiko for introduksjoner av arter med fartøy. Andre vektorer for introduksjon av fremmede marine arter blir også evaluert, slik som den økende mengden med marint søppel som kan føre med seg begroingsorganismer, egg og larver til nye områder. En analyse av historiske data for etablerte fremmede marine arter viser at kysten av Sør-Norge er særlig mottagelig for fremmede arter. Dette mønsteret reflekter ikke bare vektorpresset, men også temperaturgradienten når man går nordover i landet.

Mulig tiltak for å begrense tilførselen av nye fremmede arter til norskekysten og tiltak mot problematiske arter blir også diskutert.

Content

1	Introduction	6
2	Mapping and monitoring	8
2.1	Red king crab (<i>Paralithodes camtschaticus</i>)	8
2.2	Snow crab (<i>Chionoecetes opilio</i>)	9
2.3	American lobster (<i>Homarus americanus</i>)	11
2.4	Pacific oyster (<i>Crassostrea gigas</i>)	12
2.5	Pink salmon (<i>Oncorhynchus gorbuscha</i>)	14
2.6	Round goby (<i>Neogobius melanostomus</i>)	16
2.7	<i>Didemnum vexillum</i>	16
2.8	<i>Agarophyton vermiculophyllum</i>	19
2.8.1	<i>Zooplankton</i>	20
2.8.2	<i>Mnemiopsis leidyi</i>	20
2.8.3	<i>Gonionemus vertens</i>	23
2.8.4	<i>Acartia tonsa</i>	24
2.9	Benthic species	24
2.10	Species in ballast water	26
2.11	Species on vessels	27
2.12	Parasites and pathogens	29
2.13	Other mapping programs	30
2.14	Summary of monitoring and mapping in Norwegian Waters	31
3	Vectors for introduction of NIMS into the Norwegian coast	33
3.1	Ballast water	33
3.2	Biofouling on vessels	35
3.3	Analyses of shipping activity in Norwegian ports	36
3.3.1	<i>Cargo</i>	37
3.3.2	<i>Tankers</i>	37
3.3.3	<i>Tugs and special crafts</i>	38
3.3.4	<i>Fishing vessels</i>	39
3.3.5	<i>Passenger vessels and leisure crafts</i>	40
3.3.6	<i>Svalbard</i>	41
3.3.7	<i>The ports behind the ports</i>	43
3.3.8	<i>Geographical distribution and origin of foreign arrivals in Norwegian Counties</i>	44
3.4	Import and sale of living seafood	58
3.5	Transport of live cleaner fish	59
3.5.1	<i>Risk factors related to translocation of live cleaner fish</i>	60
3.6	Trade of species for saltwater aquaria	61
3.7	Species transported by floating debris	61
3.8	Transport of species with fishing gear	62
3.9	Intentional introduction of species and import of species for aquaculture	63
4	Ecological impact assessment of vectors, species, and biogeography	64
4.1	Alien species currently reproducing in Norwegian waters	64
4.2	Door knocker species to Norwegian waters	71
5	Management of dispersal vectors and high impact species	78
5.1	Measures for prevention of introduction and dispersal of NIMS	78

5.2	Measures for eradication of problematic NIMS	79
5.2.1	<i>Pathogens</i>	79
5.2.2	<i>Pacific oysters</i>	79
5.2.3	<i>Pink salmon</i>	79
5.2.4	<i>Didemnum vexillum</i>	80
5.2.5	<i>Round goby</i>	80
6	References	82
7	Appendix 1	92
8	Appendix 2	114

1 - Introduction

Introduced non-indigenous marine species (NIMS) can cause large ecological and economic consequences in their new environments, and monitoring their presence and distribution is essential for mitigating potential negative consequences. Currently there are only 42 recorded NIMS in Norwegian waters. This is in sharp contrast to the situation further south in Europe and particularly in the Mediterranean, where 900 NIMS are established. Sea temperature is in general the most important factor regulating the distribution of marine species. In Norway, winter temperature is often too low for species present further south in Europe to survive, or the summer temperature is too low for the species to reproduce. This is also the case for introduced species and is most likely reflected in the low number of NIMS in Norwegian waters.

The establishment and dispersal potential of NIMS in Norway and the potential ecological impact on native species and ecosystems is assessed by expert groups arranged by the Norwegian Biodiversity Information Centre and revised with regular intervals. A risk assessment was also performed for several door knocker species (defined as alien species which has not yet established in Norway but is likely to do so within 50 years) in the last revision in 2018. For the new revision in 2023, a more extensive assessment of potential door knocker species is performed through horizon scanning of species listed for the European region.

Monitoring of established populations of NIMS can have different purposes, such as stock assessment of economical valuable NIMS, surveillance of population and dispersal development of troublesome species, screening of pathogens in organisms and repeated investigations of sites with high risk for new introductions. A general mapping of established species gives good information on their dispersal, abundance and is useful for assessing their risk to native ecosystems. General mapping, which also has the purpose of detecting newly arrived species, is useful for early warning of species with a high risk of ecological impact but also for providing an inventory of NIMS present in the country and their distribution area.

Our neighbouring countries have several introduced species which have not yet established in Norway, such as round goby (*Neogobius melanostomus*), Japanese shore crab (*Hemigrapsus sanguineus*) and Harris mud crab (*Rhithropanopeus harrisi*). Alien species occurring in Denmark and Sweden can arrive in Norway unaided through movement of adults or propagule dispersal by sea currents. Introduction of a new species can be a stochastic event where one foreign vessel that arrives to a port this year brings a clonal species which establishes instantly. In most cases, a high vector pressure into an area is more likely to bring new marine species and there is probably a need for multiple introductions for a species to successfully establish, in particular in species with sexual reproduction. Assessing vector pressure on a temporal and spatial scale is thus a meaningful and relevant exercise.

Worldwide there is an increasing focus on the problem with species on the move and measures to prevent introductions of new marine species. With the implementation of ballast water treatment, an important vector for marine introductions is minimized, but still the vessels that invested in expensive treatment systems may anchor up beside a heavily fouled vessel with alien biota. In Norway, several measures to prevent introduction of new marine species are in place, and vectors such as import of alien species for aquaculture and living seafood are regulated, while vectors such as transport of living cleaner fish and floating debris still represent a risk of introducing new marine species.

In 2021 the Institute of Marine Research (IMR) in corporation with the Norwegian Institute for Nature Research (NINA) received an assignment from the Norwegian Environment Agency. Bergen Port Authorities has contributed with analysis of the cruise traffic. The assignment should focus on elucidating the current mapping

and monitoring activity on alien marine species, assess vectors for dispersal, present monitoring methods and develop a proposal for a national monitoring plan for the marine environment. This report is the first of two reports to complete the assignment. This report gives an overview of existing monitoring and mapping activities on marine alien species in Norwegian waters. Moreover, an analysis of the vectors for such introductions are given. The second report (in Norwegian) will focus on methods for mapping and monitoring and gives a proposal for a national monitoring plan.

2 - Mapping and monitoring

2.1 - Red king crab (*Paralithodes camtschaticus*)

The red king crab is an introduced species to the Barents Sea (Figure 1). It is native to the North Pacific Ocean as Bering Sea, Gulf of Alaska, Sea of Okhotsk and Japan. Russian scientists released red king crabs near Murmansk with the goal to create a sustainable fishery to increase the living standards for the local human population (Orlov & Ivanov 1978, Jørstad et al. 2002). The first transfers were made already in the 1930's but presumably none survived the journey to actually be released. In 1961, 1.5 million larvae were released in the Bay of Murmansk and over the coming years until 1969 10 000 juveniles and 2 609 adults were released in the same area. The released animals originated mainly from Peter the Great Bay in the Sea of Japan, as well as off southwestern coast of Kamchatka in the Sea of Okhotsk. In 1977 to 1978, further 1 200 adults of red king crab originating from the Far East (no further details given) were released in the Murmansk area (Kuzmin & Olsen 1994). It was estimated to take from 10 to 15 years before a population would be fully acclimatized to the new area in the Barents Sea (Orlov & Karpevich 1965).

The first few crabs were captured in the Murmansk area in 1974, and in 1976 on the Norwegian side (Kuzmin et al. 1996). A ban on fishing red king crab in the entire Barents Sea was agreed upon during the 1976-1977 negotiations between Norway and the Soviet Union (Sundet & Hoel 2016). In 1992, the red king crab was numerous in Varangerfjorden and had become a pest to the local fishermen. By November the same year, the Joint Russian-Norwegian Fisheries Commission requested both countries to intensify and co-ordinate further investigations (Kuzmin et al. 1996). In 1994, a management decision was agreed upon with a male-only fishery, as well a minimum legal size (Sundet 2014). The red king crab has since increased in population size and expanded westward to Norwegian coastal areas, from the Russian border to Hammerfest in west (Hjelset 2012, Sundet & Hoel 2016).

The red king crab is among one of the largest arthropods of the world and the population in the Barents Sea is thus also a valuable commercial species for Russia and Norway. Norway started a small experimental fishery in 1994 with 37 tonnes, but as the crab expanded its population size and distribution range the quotas increased (Sundet 2014). A commercial fishery commenced in 2003 and the quotas was set to 840 tonnes. Currently, in the Russian Economic zone the fishery is quota regulated with about 9 000 to 10 000 tonnes annually. Norway has implemented a dual management system. East of 26°E the annual goal is 1 000-2 000 tonnes, and west of 26°E the fishery is free on all sizes aiming to reduce further spread of the red king crab.

Red king crab has a larval phase consisting of four zoeal and one post larval stage before metamorphosis to juvenile crab (Stevens 2014). The transitional stage is usually referred to as glaucothoe, like megalops in brachyuran crabs. With the increasing presence of an introduced species the questions of biological characteristics, as well as ecological impact have always been given attention. Sundet (2014) made an overview of the new population of red king crab in the Barents Sea with regards to several biological questions as e.g., growth, reproduction, diet and seasonal movement. Hjelset (2012) investigated life-history parameters of the females with special focus on temporal and spatial spreading in three Norwegian fjords. As well, Oug et al. (2011) documented effects on the benthic fauna due to presence of red king crab in the Varangerfjord, an area with very high densities.

The red king crab is monitored by the Institute of Marine Research (IMR) through two surveys conducted annually: one in early summer with pots in the free fishing area, west of 26°E, and one in autumn in the commercial fishery area. (Figure 1, Figure 3).



Figure 1. (left) Red king crab (*Paralithodes camtschaticus*). Photo: Erling Svensen. (right) Monitoring of red king crab in Norway, red area shows where monitoring is taking place.

2.2 - Snow crab (*Chionoecetes opilio*)

The snow crab (Figure 2) is a subarctic species native to North Pacific, Chukchi and Beaufort Seas in the arctic and in the Northwest Atlantic from Gulf of Maine to Labrador and in the St. Lawrence Gulf as well as on the east coast of Greenland (Slizkin 1982, Williams 1984). Thus, snow crab has not previously been described as naturally occurring in the Barents Sea. However, in 1996 five individuals (four adult males and one adult female) were captured as bycatch on various vessels from May to November (Kuzmin et al. 1999). Since the first findings in 1996, snow crab is now established as a self-sustaining population and further expanded its distribution range and population size in the Barents Sea (Kuzmin 2000, Alvsvåg et al. 2009, Agnalt et al. 2011). A small-scale fishery commenced in 2012 with 2.5 tonnes, but annual harvest has increased along with increasing population size and was 20 000 tonnes in 2021 (Norges Råfisklag <https://www.rafisklaget.no/>). Since 2017, only vessels from Norway and Russia are participating.

Snow crab is a stenothermic species found at bottom temperatures below 5°C, at depths from 50 to 400 m, often linked with muddy substrate (e.g. Dionne et al. 2003). Snow crab in the Barents Sea are found at depths and temperatures like e.g. in Northwest Atlantic and North Pacific (Alvsvåg et al. 2009). Juvenile snow crabs are more stenothermic, i.e. less tolerant to slightly colder and slightly warmer temperatures than adult crabs (Alvsvåg et al. 2009, Dionne et al. 2003). The larval phase consists of three stages, zoea I, zoea II and megalope, lasting from three to four months depending on temperature (Ouellet & Sainte-Marie 2018). It is unclear how the snow crab entered the Barents Sea ecosystem. Kuzmin et al. (1999) introduced the theory of ballast water as a pathway, through tankers coming from the northwest Atlantic. Crab species like e.g. snow crab that have a long pelagic larval phase can survive intercontinental voyages in ballast water. One such example of accidental introduction is the green crab *Carcinus maenas* (e.g. Darling et al. 2008). The ballast water hypothesis implies an introduction of relatively few individuals possibly creating a genetic founder effect or bottleneck. There is no genetic evidence of such concerning the snow crab in the Barents Sea (Dahle et al. submitted). Trans-arctic interchange has also been suggested as a possible pathway (Agnalt et al. 2011). A third theory is natural expansion by individuals moving from established populations along the north coastline of North

Russia/Siberian coast (Dahle et al. submitted).

The population of snow crab in the Barents Sea is monitored by IMR through a joint Norwegian Russian Ecosystem survey, taking place during both winters i.e., January/February and summers i.e., August/September (Figure 3). In addition, a dedicated snow crab survey has taken place since 2018, run during June (Hjelset, pers. comm.). IMR has currently one internal funded project monitoring the snow crab in the Barents Sea that covers the cost of the pot survey (Figure 2). In addition, the project Snowman (project no. 14862; 2018-2021) was initiated to answer several questions relating to the biology of a new species which may differ in the non-native area compared to regions where the species has been established for longer periods (e.g., Brockerhoff & McLay 2011). In SnowMan, focus was set on genetics to elucidate the donor population of the snow crab in the Barents Sea (Dahle et al. submitted), biological characteristics as fecundity (Danielsen et al. 2019), size at maturity, various disease/ectoparasites aspects as e.g. the bitter crab disease being common in other snow crab populations (Nunkoo et al. submitted), detection of larvae in western part of the Barents Sea (Hjelset et al. 2021) and if this can contribute to explain further spreading in the western direction (Huserbråten et al. in prep). Further, what does snow crab consume in the Norwegian zone (Sundet et al. in prep) and what is the impact of a large snow crab population on the benthic production (Holte et al. in prep), as well as impact on the ecosystem level (Hansen et al. in prep). The Northeast Atlantic cod do prey on snow crab (Holt et al. 2021) and can prey-predator be used as a stock assessment index.

As snow crab is a high-priced product, the established population in the Barents Sea has been expected to accommodate a significant fishery. Snow crab was in 2015 defined as a sedentary species i.e. Russian and Norwegian property on their respective portions of the continental shelf. These rights extend beyond the 200 nautical miles of both the Russian and the Norwegian EEZ. This has been disputed and consequently the fishing right has been a political issue for several years (e.g., Hansen 2016, Østhagen & Raspotnik 2018, Kaiser et al. 2018).



Figure 2. (left) Snow crab (*Chionoecetes opilio*). Photo: Ann Merete Hjelset. (right) Monitoring of snow crab in Norway, red area shows where monitoring is taking place.



Figure 3. (left) Pot-fishing of snow crab. Photo: Ann-Merete Hjelseth. (right) Monitoring area of snow crab through the Ecosystem Survey Monitoring Program 2004-2021. Red area indicates area which is investigated.

2.3 - American lobster (*Homarus americanus*)

American lobster (Figure 4) is native to the North-Eastern Atlantic, from Cape Hatteras (USA) in south to Newfoundland (Canada) in north. It is a coastal species, captured mainly by pots. The fishery has increased from 36 851 tonnes in 1980 to 161 011 tonnes in 2019 (FAO Statistics). Consequently, export has also increased accordingly. In 2015, approximately 75 500 tonnes of *H. americanus* was exported from Canada and USA (FAO Statistics). Barrett et al. (2020) reports that in 2015 UK imported 1 700 tonnes live specimens. Europe has traditionally been the important market, but the last decade China and other Asian countries have increased in importance. Import of live animals poses a risk when accidentally or intentionally released into the environment in the area/location/country of import. The American lobster is geographically separate from the European lobster by the deep Atlantic Ocean.

Import of live specimens has led to escaped individuals and intentional release into the European marine environment. Findings of live American lobster have been confirmed in various European countries like Denmark, Sweden, UK, Ireland, Croatia i.e., the Adriatic Sea and Norway (Jørstad et al. 2011, Stebbing et al. 2012, Øresland et al. 2017, Barrett et al. 2020, Pavičić et al. 2020). In UK, a total of 162 individuals have been identified based on morphological characteristics such as spine below the rostrum and colouration, over the period from 2012 to 2018 (Barrett et al. 2020). In UK, the majority can be traced back to an event of which 361 American lobsters were released in the English Channel 15th June 2015, as part of a Buddhist ritual (Barrett et al. 2020). Also, 35 non-native Dungeness crab (*Metacarcinus magister*) were released in the same event. In Norway, a total of 35 specimens have been identified genetically as American lobster over the period from 2000 to 2017 (Agnalt pers. comm.). Crossbreeding with local native European lobster have been found in five American females in Norway, two in Sweden and one in UK (Agnalt pers. comm., Barrett et al. 2020).



Figure 4. American lobster (*Homarus americanus*) with hybrid eggs. Photo: Beate Hoddevik.

In 2016, it became illegal to import live American lobster to Norway. The same year the Government of Sweden proposed an EU-wide ban on importing live American lobster. Hybridization with the native European lobsters were raised as concerns and possible negative impacts on other native species such as edible crabs (*Cancer pagurus*). In addition, possible hitchhikers or biofouling on the American lobsters included several other non-native species (nematodes, copepods, barnacles and polychaetes). However, EU did not approve the suggestion of the Swedish government.

In Norway, there are no current monitoring program for American lobster. Detection is solely based on observant fishermen (recreational and commercial).

2.4 - Pacific oyster (*Crassostrea gigas*)

The Pacific oyster, *Crassostrea gigas* (Figure 5), was introduced to Scandinavia and farmed at several sites in the 1980's and early 1990's. Despite the farming activities, feral populations did not establish. A bio-invasion of Pacific oyster however commenced in Scandinavia in 2007 (Wrangle et al. 2010), and the oyster is now established along most of the Scandinavian coastline, approximately north to Bergen, Norway. Genetic studies (Faust et al. 2017), combined with the pattern of spreading (Wrangle et al. 2010), suggest that most of the oysters spreading in Scandinavia have the same origin, and spread northwards in a "stepping stone pattern". The Scandinavian populations may already have adapted to local conditions (Sussarellu et al. 2015), and reproduction in new areas may be facilitated by warm summers. Larvae are spread with the water currents and settle on suitable substrates.

The bio-invasion has been followed since 2007 and the colonized habitats have been assessed (Mortensen et al. 2017). Reise et al. (2006) described the invasion in four phases. The invasive species arrive to the new habitat (Phase 1) and start to establish a population at low density without changing the habitat (Phase 2). As the population starts to expand it increases in density and may change the habitat (Phase 3). Once established in the habitat other species as predators, competitors and diseases/parasites will reduce the density in an adjustment phase (Phase 4), potentially reducing, or at least altering, the effects of the non-native species on the ecosystem.

In most of its range in Scandinavia, populations have gone through several bottlenecks, like mortalities during cold winters (Strand et al. 2012) and summer mortalities after hot summers (Mortensen et al. 2016). This may correspond to Phase 4. In the outer edges of its range; along the Norwegian west coast, it may still be in Phase 1.

In accordance with the proceeding of the invasion process into phases 3 and 4, we have moved from mapping and studies of the mechanisms of the invasion, to the establishment of a monitoring based on a common Scandinavian model. Along the western coast of Norway, the Pacific oyster has not yet established in all available habitats and is considered in Phase 1 and 2. A mapping is now performed to clarify the process of spreading from Rogaland and northwards. Data from the mapping will be prepared for publication and presented in 2022 (Figure 5).

The Scandinavian studies on the establishment, spreading and mitigation of Pacific oysters has, since the bio-invasion commenced in 2007, been organized as a collaborative network project, including scientists and managers from Norway, Sweden and Denmark (see Mortensen et al. 2019).

To help with the distribution mapping, the Institute of Marine Research (IMR) has invited the public to submit information, photos, coordinates etc, on occurrences of Pacific oysters. A [video](#) showing the differences between flat oysters and Pacific oysters and typical characteristics has also been published to aid the identification. Information from the public has been evaluated by experts and all credible data included in distribution maps (publication in prep). The Scandinavian network has been funded by The Nordic Council of Ministers in three periods since 2011, ending in 2021. Participants from each country have been funded through national projects linked up in the network.



Figure 5. Pacific oyster (*Crassostrea gigas*). Photo: Vivian Husa. (right) Monitoring of pacific oysters in Norway, red area shows where monitoring is taking place.

2.5 - Pink salmon (*Oncorhynchus gorbuscha*)

The anadromous pink salmon (Figure 6) is native to the Northern Pacific. It has been released repeatedly in Russia, and in recent years it has occurred in high densities in Norway. Surveillance of the distribution and abundance of invasive pink salmon in Norway started in 2017 (Berntsen et al. 2020). Surveillance of pink salmon is mainly conducted through registration of catches in sportfishing or targeted removal fishing in rivers and catches in bag- or bend-nets in the sea (along the coast) by fishermen. In addition, occurrence and abundance of pink salmon is registered by drift counting surveys, camera surveillance and by environmental DNA (eDNA) surveys in selected rivers (Figure 6). Pink salmon occur in high densities particularly in Northern Norway (Figure 7). The surveillance activity is organized by local organisations and commercial companies in association with the County governor. Catches of pink salmon at sea are also registered through surveys by the Institute of Marine Research.



Figure 6. (left) Pink salmon. Photo: Christine Fagerbakke. (right) Monitoring of pink salmon in Norway, red area shows where monitoring is taking place.




Figure 7. Map showing the total number of registered pink salmon, *Oncorhynchus gorbuscha*, in Norwegian rivers in 2017 and 2019. Fish registered are either caught or observed, by several methods. Rivers included are from the NVE database, while some small rivers are not included (from Berntsen et al. 2020).

The Norwegian Institute for Nature Research (NINA) works as an advisor for the Norwegian environment agency regarding monitoring (registration of catches) and biological sampling of pink salmon. NINA also gathers all available data on pink salmon in a common database and report analyses of these data in NINA-reports, which are available at <https://www.nina.no/pukkellaks>. NINA was also the national coordinator for a cross-Atlantic citizen science project using eDNA for detection of pink salmon in rivers in 2019 (www.1000rivers.net).

2.6 - Round goby (*Neogobius melanostomus*)

The round goby (Figure 8) is an invasive fish from the Ponto-Caspian region, which is spread widely in both Eurasia and North America (Kornis et al. 2012, Forsgren & Florin 2018). It is a door knocker species for Norway associated with severe impact (Norwegian Biodiversity Information Centre 2018). There is currently no existing monitoring program for early detection of the round goby. There are, however, a couple of ongoing projects focusing on the species. NINA is currently analysing eDNA samples collected from Trondheim harbour and the Oslofjord area in 2021, with samples from Halden, Hvaler, Fredrikstad, Oslo harbour, Drammen and Sandefjord. The Institute for Marine Research in cooperation with NORCE is supervising a MSc-project in 2021 that is investigating possible occurrence of the species with eDNA, fishing by hook and minnow traps from Gothenburg in Sweden into the Oslofjord. So far, the species has not been detected in Norway (Figure 8).



Figure 8. (left) Round goby caught by hook in Gøteborg. Photo: Vivian Husa. (right) Target areas (red) of ongoing projects investigating possible introduction of round goby in Norway.

2.7 - *Didemnum vexillum*

The invasive colonial ascidian *Didemnum vexillum* (Figure 9) was recorded near Stavanger for the first time in Norwegian waters in November 2020. The species is one of the world's worst alien species as it grows quickly, cover and hampers marine habitats from 0-65 meters depth (McKenzie et al. 2017 and references therein) It is also expected that it will have economic impact as it grows willingly on oyster and mussel cultures, kelp and finfish farms. Moreover, it can cause the shipping industry considerable cost for cleaning of vessels going out from infected areas. The species have temperature tolerance limits between -2 and 24 C°, (McKenzie et al.

2017 and references therein) which means it has the potential to establish along the entire coastline of Norway.

A monitoring project of the growth potential of *D. vexillum* colonies during the year was set up in 2020 in the Stavanger area in corporation with Stavanger diving club, the Museum of Stavanger, and Stavanger Municipality. Preliminary results from this project show that the colonies were shrinking a bit in March-April when the water temperatures were below 4 C°, but a remarkable growth of colonies continued during the summer/autumn season 2021.

The species grows best in salinities above 25 psu and dies under 20 psu (McKenzie et al. 2017 and references therein), and its distribution will likely be restricted in inner parts of Norwegian fjords with low salinity. During August 2021 a survey of 70 stations in Rogaland was performed by diving, video recordings and investigations of structures in ports and marinas. The survey showed that the species was very abundant in Engøysundet (Stavanger) where it was first recorded. The species covered up to 50 % of the seafloor with highest abundance close to a moored pram, where probably a vessel with *D. vexillum* on the hull has been situated and dripped to the seafloor (Figure 10). 50-60 stations in Rogaland were investigated in this survey, with no more records of *D. vexillum*.

In September 2021 we received reports from divers in Haugesund (60 km further north in Rogaland County) with new records of the species. In this area the species is more widespread, with a dense cover in Karmsundet and around Vibrandsøy (Figure 10). Data from the Haugesund area is obtained from trained divers from Sletta Dykkeklubb. Later in September, the species was discovered from an oil rig in a commercial port at Askøy north of Bergen (Vestland county) (Figure 10). *D. vexillum* was growing on the seafloor at the port and has spread to three more locations outside the port. A survey in Bergen harbour in October yielded no records of *D. vexillum*. In December 2021 a diver reported several colonies (Erling Svensen pers. comm.) at three sites in Egersund approximately 80 km south of Stavanger. This makes it clear that the species is spreading fast between busy port cities.



Figure 9. *Didemnum vexillum*. Photo: Erling Svensen.

These surveys were performed as a part of an internal IMR project funded by the Ministry of Trade, Industry and Fisheries. A monitoring project was initiated by the Norwegian Environment Agency in November 2021 to use e-DNA for mapping of *D. vexillum* in ports from Stavanger to Bergen. This mapping is performed by NINA in corporation with IMR. The results from the e-DNA mapping showed strong signals at all sites where *D. vexillum* has previously been recorded and some sites nearby, where the species occurrence also was confirmed recently by visual investigations. There were no e-DNA signals between Bergen and Haugesund, but a strong signal at several sites in Gulen municipality north of Bergen. *D. vexillum* was recorded at one of these sites, a lay-up place for oil rigs, in February 2022. The study also showed some uncertain or unclear e-DNA signal at several sites, which will be checked up by visual surveys forthcoming (Fossøy et al. 2022, unpublished data IMR).




Figure 10. Distribution of *Didemnum vexillum* in Norway and year of first record in specific area (Maps from the Norwegian Biodiversity Center).

2.8 - *Agarophyton vermiculophyllum*

The red algae *Agarophyton vermiculophyllum* (Figure 11) was recorded for the first time at the Norwegian coast near Tønsberg (Vestfold & Telemark County) in 2012. As this species has become very abundant in several countries, an annual mapping program in inner and outer Oslofjord was performed in the period 2013 to 2019 (Figure 11). The mapping has been limited, and only a part of the coastline has been examined each year, but some stations have been investigated several times. As *A. vermiculophyllum* thrive best in wave protected, shallow and muddy habitats, the mapping have been restricted to such areas. Abundance of the species has been done with observations (beach survey) and collection of material with a grapnel. The abundance of *A. vermiculophyllum* in seagrass meadows (*Zostera marina*) was studied in three areas in Oslofjord in 2020 and showed only low abundance in this habitat in Viksfjorden, Tjøme and inner Oslofjord (Sundal-Joys 2021).



Figure 11. (left) *Agarophyton vermiculophyllum*. Photo: Rudolf Svensen. (right) Monitoring of *A. vermiculophyllum* in Norway, red area shows where monitoring is taking place.

2.8.1 - Zooplankton

Zooplankton are animals that live all or part of their life drifting in the water column. Zooplankton includes a wide range of animals, such as copepods, cladoceran, jellyfish and the larvae of benthic invertebrates. Ballast water is one of the primary transport vectors for the transfer and introduction of non-indigenous zooplankton, while ocean currents play an important role for the secondary spreading.

The institute of Marine Research operates several zooplankton monitoring programs in all Norwegian seas, including open ocean as well as coastal waters. The IMR zooplankton monitoring focuses on meso-zooplankton (size 180 μm -20 mm) and the sampling methods are usually targeting crustacean plankton. Although the monitoring aims to include all taxa, there is no dedicated monitoring program specifically addressing alien zooplankton species. Information on occurrences of alien zooplankton in Norwegian waters are derived from various sources, often as qualitative data. A few examples of non-indigenous marine zooplankton species in Norway are mentioned below.

2.8.2 - *Mnemiopsis leidyi*

Mnemiopsis leidyi is a lobate ctenophore (comb jelly) native to the east coast of North and South America (Figure 12). The species was introduced with ballast water to Europe (the Black Sea) in the early 80's. The first observation of *M. leidyi* in Norwegian waters was made in Oslofjord, November 2005 (Olivera 2007). During 2005 – 2007 the species was recorded in several areas of northern Europe, including southern North Sea, Kattegat, and the Baltic (Javidpour et al. 2006, Hansson 2006, Boersma et al. 2007). The introduction to Northern Europe in the 2000's probably occurred as an independent introduction, directly originating from the northeast coast of the USA, probably with ballast water (Costello et al. 2012).

Since 2005, the species has occurred in high densities in Norwegian coastal waters during August-December, from Oslofjord to Trondheimsfjorden. The exception is the years 2011-2013 when *M. leidyi* was absent, probably due to low winter temperatures. Binndalsfjorden (approx. 65.2° N) is the northernmost observation made so far by *M. leidyi* on a global basis.

M. leidyi is a hermaphrodite (bisexual) with high reproductive capacity and the ability to self-fertilization. *M. leidyi* is highly adaptable and tolerates a wide range of temperatures and salinities. The species may occur in temperatures from -0.7 to 35 °C and in salinities between 3.4-70 ppt (Hansson 2006, Costello et al. 2012). Furthermore, the species survives food shortages for as long as three weeks or more. These traits enable *M. leidyi* to survive long transport in ballast water, and quickly colonize new areas.

M. leidyi is a carnivore with a broad food spectrum, including small zooplankton, copepods, fish eggs and larvae (Næss 2015). The species have a high predation rate and is able to eat up to ten times its body weight in food per day. In areas with dense populations of this ctenophore, it may locally have strong effect on zooplankton populations which can and affect several trophic levels (cascade effect, Granhag et al. 2011, Tiselius and Møller 2017).

M. leidyi spends the entire life cycle in the free, marine water masses (holoplanktonic) and spreads rapidly with ocean currents. The coastal current is an important dispersal route for the species, and its distribution in Norwegian waters is linked to coastal areas and within fjords. The temperatures and salinities along the entire Norwegian coast are within the species' tolerance level (Hosia & Falkenhaus 2015), and the species has the potential to overwinter by seeking deeper (and warmer) water layers throughout the winter. It is expected that the species will colonize areas further north along the coast, due to future higher temperatures. Reproductive stages of *M. leidyi* have been observed in coastal waters in August-September. However, *M. leidyi* has so far not been recorded in Norwegian waters during the winter. This indicates that the occurrence in Norwegian sea areas is driven by so-called «Source-sink dynamics», where the population is maintained by the annual supply of individuals from overwintering areas in southern North Sea (Jaspers et al. 2006, Costello et al. 2012, Hosia and Falkenhaus 2015).

There is no coordinated monitoring program, targeting the distribution and spreading of *M. leidyi* in Norway. Gelatinous plankton is recorded as by-catch in the standard zooplankton monitoring carried out by the Institute of Marine Research. However, the methods used (plankton nets) is targeting crustacean plankton and will not give reliable data on delicate gelatinous plankton. Nevertheless, records of *M. leidyi* from plankton nets and the beach seine programme (Skagerrak), have provided qualitative information on distributions as well as seasonal and interannual variations of this species (Figure 12).




Figure 12. (left) *Mnemiopsis leidyi*. Photo: Erling Svensen. (right) Mapping area (red) of distribution of *M. leidyi* in Norway.

Observations made by the public are an important supplement to the information on distributions and blooms of *M. leidyi* in coastal waters. Information on the species, and how to distinguish *M. leidyi* from the native *Bolinopsis infundibulum* is posted on the IMR website. The IMR receives numerous observations of *M. leidyi* from the public each year by email. Since 2019 a new platform for Citizen Science has been used (<https://dugnadforhavet.no/>) with a separate section on gelatinous plankton. This application has provided georeferenced records of the species, particularly during bloom events. In 2021 more than 50 observations of *M. leidyi* were received from the public via “Dugnad for Havet” (Figure 13).

The research project “Distribution and ecosystem impacts of the invasive ctenophore *Mnemiopsis leidyi* in Norwegian waters” was funded by the Norwegian Research Council 2009-2013.




Figure 13. Observations obtained from the Citizen science application "Dugnad for havet" in 2021. a) *Mnemiopsis leidyi*, b) *Gonionemus vertens*.

2.8.3 - *Gonionemus vertens*

Gonionemus vertens is a small hydromedusa (20-25 mm in diameter), native to the northwest Pacific Ocean (Figure 14). The species is often associated with eelgrass beds and macro algae in sheltered, coastal areas. The medusae is "semi-planktonic" and attaches to eelgrass and seaweed during the day using specialized "adhesive plates" on the tentacles. *G. vertens* was observed in the North Sea in 1913 and then spread throughout the North Sea area until the middle of the 20th century. The first observation of *G. vertens* in Norway is from 1921, at Drøbak in the Oslofjord (Kramp 1922). Later, two individuals were found in Hardangerfjorden in 1958 (Tambs-Lyche 1964) and Trondheimsfjorden 1969 (Gulliksen 1971). The northernmost observation was made in 2020, at Andenes (approx. 69.3° N). Since 2016 the number observations of *G. vertens* have increased, especially in southwestern Norway (Rogaland-Agder), indicating a possible new introduction to Europe. There have also been several reports on severe stings to humans in Norway. Regional variations in burning effect (toxicity) indicate that there may be several genetic lineages of the species. The recent records of *G. vertens* in both Sweden and Norway probably belong to a more toxic phenotype, compared to the former (historical) records of the species (Govindarajan et al. 2019). There is great uncertainty about how *G. vertens* spreads to new areas. The species is probably transported on ships' hulls at the polyp stage and by ballast water and currents (medusae stage and eggs). It is also believed that polyps of the species were introduced to Europe in connection with the import of oysters from Japan and the United States.

There is no coordinated monitoring program, targeting the distribution and spreading of *G. vertens*. The preferred habitat for this species (shallow areas, within the algae belt) is not covered by standard zooplankton monitoring. Data on occurrences of *G. vertens* is based on observations made by the public. The medusae is small and is easily overlooked unless people get stung. Since 2019 observations of *G. vertens* submitted

through the IMR platform for Citizen Science (<https://dugnadforhavet.no/>) has provided valuable georeferenced records.



Figure 14. (left) *Gonionemus vertens*. Photo: Erling Svensen. (right) *Acartia sp.* Photo: IMR.

2.8.4 - *Acartia tonsa*

Acartia tonsa is a small planktonic copepod (Figure 14), native to American and Indo–Pacific waters (Leppäkoski & Olenin, 2000), but has been spread worldwide and is now regarded as cosmopolitan. In Europe it was first discovered in 1916, and in Sweden it was registered in 1934. The first Norwegian record of *A. tonsa* was made in 2012, in a land locked fjord in Southern Norway (Landvikvannet, Haraldstad et al. 2013) and the species has recently been registered in Skagerrak coastal waters (Moseid et al. 2021).

A. tonsa is a coastal and estuarine species that occurs in brackish water areas with reduced salinity (15-20 psu) and well adapted for highly eutrophic waters. The species requires temperatures of at least 10 oC for reproduction. *A. tonsa* produces benthic diapause eggs, which survive the winter and hatch in the spring when temperature reaches 10 °C. The ability to produce highly resistant diapause eggs has probably played an important role for its wide distribution. Ballast water seems to be the most likely means of transferring copepodites or resting eggs of this species. *A. tonsa* is planktonic, and therefore spreads rapidly with the coastal current.

Acartia sp is not identified to species in the IMR standard monitoring programs. It is probable that the species is present in several fjords and estuaries. However, brackish water areas are generally a poorly studied habitat type in Norway and the distribution of *A. tonsa* along the Norwegian coast is poorly known. *A. tonsa* is morphologically very similar to native *Acartia* species, and it is time-consuming and demands taxonomic expertise to identify and monitor this species by traditional morphological techniques. Molecular based species-specific assays have been developed and proven to be a powerful tool to confirm the presence of *A. tonsa* in mixed plankton samples (Moseid et al. 2021).

2.9 - Benthic species

A mapping program for benthic introduced species in marinas along the coast were initiated in 2010 and is still

ongoing (Figure 15). This investigation is inspired by the methods used in Rapid Coastal Survey RCS (Minchin 2007) and focuses on targeted established NIMS and door knocker species. The general idea is to use a cost and time efficient method to provide a general picture of the distribution of species. The methods have developed during the years and now includes beach survey, snorkelling and diving at selected localities. The field work is usually performed during five days in summer with several teams examining 50-70 stations altogether in a selected area of the coast. Approximately 450 marinas and beaches along a coastline of approximately 1750 km have been covered in this survey, some stations twice (Husa et al. 2012a, b, 2013). 18 established NIMS were recorded in this project, among those the four recent newcomers *Schizoporella japonica* (2014), *Agarophyton vermiculophyllum* (2012), *Grateloupia turuturu* (2018) (Figure 15) and *Didemnum vexillum* (2020).



Figure 15. (left) *Grateloupia turuturu* at Ula harbour. Photo: Rudolf Svensen. (right) Mapping area (red) of distribution of targeted door knockers and established NIMS in marinas in the period 2010-2019.

We have also tested out the efficiency of other methods such as dredging in the subtidal, fishing nets, traps and crab pots during the project. We also tested out a complete harbour inventory in the port of Narvik in 2012 (Husa et al. 2014), where all groups of taxa were collected by several methods and all species identified to lowest possible taxonomic level.

In 2016 we tried out a citizen science project where volunteers and diver club were trained to identify introduced

marine species. The success of this project was variable, but some divers and volunteers kept on looking for introduced species and reports regularly, for example new records of *D. vexillum*. An easy way to report introduced species by geotagged photos on mobile application was developed in connection with this project: <https://dugnadforhavet.no/>. Three records of *Hemigrapsus takanoi*, which has not been spotted in Norway before, was reported in the application in 2019-2021 from the Oslofjord area.

All records from these projects are reported in the Norwegian Biodiversity Centre and gives valuable information on distribution of species (Figure 16). These projects were funded by the Norwegian Environment Agency in the period 2010-2014 and since then through internal projects at IMR funded by the Ministry of Trade, Industry and Fisheries.




Figure 16. *Caprella mutica*. Photo: Rudolf Svensen. Map showing records of *C. mutica* along the coast of southern Norway (Norwegian Biodiversity Centre).

2.10 - Species in ballast water

Ocean going vessels need ballast to maintain stability, regulate buoyancy and manoeuvrability when sailing with limited or no cargo on board. Originally this was achieved by solid ballast, typically stones, soil and sand, and we do in fact have numerous examples of land plants translocated by this means

<https://no.wikipedia.org/wiki/Ballastplanter>. With the advent of water as ballast in vessels, aquatic organisms were given opportunities to hitchhike beyond their natural boundaries. The problem was addressed in a seminal paper by Jim Carlton in 1985 (Carlton 1985). Several dramatic events linked to ballast water, like HABs (harmful algal blooms), (Hallegraeff 1993) and outbreaks of Cholera epidemics in the 1990's, (Takehashi et al. 2007) fuelled a substantial research effort into the field in the following decades. The research results, and subsequent knowledge transfer to the political and managerial sectors, were acknowledged in the UN Conference on

Environment and Development (UNCED), held in Rio de Janeiro in 1992. The UN IMO (International Maritime Organization) initiated negotiations to consider the possibilities of developing an internationally binding instrument to address the transfer of harmful aquatic organisms and pathogens in ships' ballast water.

Ballast tanks are typically dark, may contain considerable amounts of sediments, and provides both "planktonic" habitat and hard-bottom substrate. Depending on sediment content, they may also provide soft-bottom substrate. Even sessile organisms may find opportunities in ballast tanks, either as adults colonizing ballast tank walls, or as the sessile organisms' usually planktonic propagule (most sessile organisms have a planktonic spreading unit (typically egg/larvae) (Kinlan & Gaines, 2003). Ballast tanks are filled and emptied by gravity, by pumps, or by a combination of these. Pumping will involve considerable pressure- and speed differences (shearing forces) that may harm larger fragile organisms. Smaller organisms and microorganisms will remain unaffected by pumping.

Since the 1980's, high number of studies around the globe have documented the large number (both of species and specimen) transported between the world's larger and smaller ports. In 14 shipping studies, Gollasch et al. (2002) surveyed more than 550 vessels entering European ports. In 1508 samples (1219 ballast water, 289 tank sediment) collected and analysed, a total of 990 different taxa were recorded. The diversity of species found included bacteria, fungi, protozoans, algae, invertebrates of different life stages including resting stages, and fishes with a body length up to 15 cm (Gollasch et al. 2002).

Arctic and Antarctic regions are not immune to the introductions of alien species (although the temperature conditions will be preventing many species from being established). In the Antarctic, live specimen of barnacles and *Mytilus galloprovincialis* have been found on vessels, and size measurements of the latter signified that the specimens had survived several visits to the Antarctic, (Lee and Chown, 2007). In the Arctic, several studies have documented transportation of live NIMS to Svalbard, (Ware et al. 2015) and in the Canadian Arctic (Goldsmit et al. 2018).

Currently the knowledge on the biogeography of bacteria is limited, but the data are clearly indicating that the "global microbiome" is geographically structured. Species and strains are to some extent "local", and the concept "alien species" therefore also is relevant for these functional groups. (Hess-Egra et al. 2019). The knowledge of the global distribution of virus is poorly studied, but newer studies have demonstrated that a) the viral content in ballast water can be high indeed, and b) several human pathogens were discovered in ballast water (Hwang et al. 2018). In Norway, only a few studies of the biota in ballast water have been made. The most comprehensive was a part of a European initiative (Gollasch et al. 2002), and a study in Svalbard (Ware et al. 2015).

2.11 - Species on vessels

The biofouling community on vessels entering the Norwegian coast from abroad has been little studied. It is difficult to get permission to inspect the hull from the shipping companies, and safety for the divers is also an issue during such operations. Most vessels are from time to time inspected by divers or ROV's, but those footages are not available to the public and often of such quality that it is difficult to distinguish species from them. In corporation with a local shipping company and an oil company IMR have inspected in total six vessels in period 2016-2019: two heavy lifting vessels and four supply vessels. Heavy lifting vessels often stays for months in a port while constructions for the oil or power industry is made ready on the shipyards, and though pose a substantial risk of getting heavily fouled. Many of these constructions are built in Korea, in waters that

have similar temperature conditions as Norwegian waters. One such vessel coming in from Korea were examined in 2018. Although the ship had been cleaned before departure from Korea, we still found several species of barnacles and goose barnacles, oysters, and other mussels in niche areas around the propeller. A similar vessel from Gothenburg were investigated the same year but had only a large number of Pacific oysters on the hull.

A supply vessel coming into the Norwegian coast after four years as a standby diving vessel at an oil field in the Gulf of Mexico, were surveyed by divers in the autumn 2016. The vessel had a moon pool and had travelled from the gulf to Norway with no bottom doors in the moon pool. The moon pool contained an almost complete ecosystem with species that are commonly associated with artificial structures in the Gulf of Mexico. In the moon pool there were swarms of blennies (*Scartella cristata*) and a yellow *Parablennius* sp. (Figure 17). Several specimens of the crab *Cronius ruber* and three species of shrimps were hiding in the holes at the sides of the moon pool. The walls were covered with sun corals, pearl-oysters and several other species of oysters and mussels, calcareous worms, sponges, ascidians, and bryozoans. A revisit at the moon pool in April the following year after a cold winter, no living animals from Mexico occurred and the pool was taken over by native ascidians and swimming crabs. One of the supply vessels coming in from Africa was completely clean, two other vessels from Chile and Brazil had remains of barnacles and empty mussels, but everything on these ships were dead because the vessels had stayed in Norwegian waters under winter conditions before the investigation. A last supply vessel was examined directly after the arrival from the Red Sea (Egyptian side) and was heavily fouled after being moored for years. This ship had a up to 10 cm thick layer of barnacles, oysters and other molluscs, ascidians, bryozoans, and sponges (Figure 18). When scraping off the fouling community it revealed that small crabs and polychaetes were living within the biofouling. We even recorded an endemic blenny (*Parablennius zvonimir*) from the Mediterranean hiding in one of the large barnacles. This hitchhiker had probably entered the ship during a few days stay in Malta on the journey to Norway.

These occasional studies are not sufficient to establish a full picture of biofouling on ships entering Norwegian waters but gives us a taste of the potential for this vector.



Figure 17. Unidentified shrimp, sun corals and Molly Miller blenny (*Scartella cristata*) collected in the moon pool of a vessel coming in to the Norwegian coast from the Mexican gulf.



Figure 18. Bilge of supply vessel coming into a Norwegian port with dense populations of mussel (*Brachidontes pharaonis*) originating from the Red Sea.

2.12 - Parasites and pathogens

Parasites and pathogens may be introduced with import or translocations of live animals for aquaculture, unintentionally via vectors or with live seafood that is re-laid in water.

Introduction and spreading of fish diseases is a well-recognized hazard, after a long period with severe disease problems and huge economical losses in the aquaculture industry. Today, the industry is restrictive, and imports are banned, to minimize the risk. A remarkable exception is the import and translocation of cleaner fish with unknown health status released in the salmon and rainbow trout net pens, and thus creating a polyculture which may lead to introductions of “new” pathogens or changes in virulence of pathogens already present in salmonids and cleaner fish species.

Spreading of pathogens is not only related to import. It is also relevant between regions in Norway. However, zoning to protect a good health status is difficult, due to the adaptation to EU legislation (former directive 2006/88, now the new Animal Health Law, regulation 2016/429). Surveillance is focused on pathogens that are notifiable, according to EU and national legislation. As a general rule, movements are allowed between areas / water compartments of equal health status or from areas of high status into areas with a lower status.

To avoid introduction of the listed / notifiable diseases, there is a need for documentation. In Norway, there is limited monitoring of fish and shellfish parasites, and pathogens present outside Norwegian waters. However, The Norwegian Food Safety Authority (NFSA) aims at obtaining an overview of the notifiable diseases. Three Surveillance programmes, funded by the NFSA are carried out:

1. The Norwegian Veterinary Institute carries out a surveillance programme for Viral haemorrhagic septicaemia virus

and infectious haematopoietic necrosis virus in farmed salmonids, pink salmon (caught in one river) and a limited number of samples from cleaner fish. The programme has a risk-based approach, and the main surveillance activity is the routine clinical inspections on sites with farmed salmonids and analyses of samples collected from diseased fish. Viral haemorrhagic septicaemia virus and infectious haematopoietic necrosis virus were not detected at any of the sites tested in the 2020.

2. The Veterinary institute also carries out a surveillance programme for Freshwater Crayfish plague. Environmental DNA (eDNA) monitoring is used for the detection of species-specific DNA from spores of *Aphanomyces astaci* directly from water filtrates. The presence/absence of eDNA from noble crayfish (*Astacus astacus*) and signal crayfish (*Pacifastacus leniusculus*) is also determined to supplement the results.
3. The Institute of Marine Research carries out a surveillance programme for bonamiosis and marteiliosis in European flat oysters, *Ostrea edulis*, and blue mussels, *Mytilus* sp. in Norway. The programme is based on a targeted surveillance combined with research activities following up mortality events. The programme combines histology and PCR-based diagnostic methods, and may thus detect other pathogens than *Bonamia* sp. and *Marteilia* sp. In the bivalve populations, and over time get an overview of the health situation of Norwegian bivalves and protect the stocks from the introduction of exotic pathogens. *Bonamia* sp. is detected in Norwegian oysters. *Marteilia refringens* Type M. I M. *pararefringens* has been detected in blue mussels from five sites. A collaboration has been established between IMR, IFREMER (France) and CEFAS (UK) to elucidate the distribution, spreading potential, life cycle and host range of M. *pararefringens*. To disseminate results and information, the programme has close contact with Scandinavian shellfish farmers. Results are published annually, through IMR report series (see Mortensen and Skår 2020).

In addition to these surveillance programmes, pathogens in wild and farmed fish and shellfish are sometimes studied in research projects and/or screening initiated by the farming industry. These activities are however not done on regular basis.

Surveillance of oysters and mussels is the only programme that may catch other pathogens than those targeted. Selected populations are monitored to detect mortality events. The general state (condition) of bivalves is noted during sampling, and histology methods are used in the health screening. This will reveal the condition of the individual bivalve, pathogenic changes in tissues and any foreign organism visible through the microscope.

The surveillance programme for bonamiosis and marteiliosis reports to the NFSA and the European reference Laboratory for mollusc diseases (EURL). This enables a dynamic collaboration on any pathogen detection or finding of abnormal tissues. The work is strengthened through collaboration with research teams at IFREMER (France) and CEFAS (UK). The work has also been linked to international research projects and a Scandinavian network used to disseminate information and exchange information and data (see below).

The surveillance programme for bonamiosis and marteiliosis is funded by NFSA, research by IMR, through funding from the Ministry NFD, and partly through research projects (EU – Vivaldi (ended 2020) and networking - Scandinavian network (Nordic Council of Ministers) (ending 2021)).

2.13 - Other mapping programs

The Norwegian Biodiversity Centre is funding several inventory projects on marine species in Norwegian waters within the program “The Norwegian Taxonomy Initiative (NTI)”. The aim of these projects is to provide inventories of all multicellular species occurring in Norway and to build validated DNA-barcode reference databases for all species. Barcoding is conducted in collaboration with the Norwegian Barcode of Life (Nor-BOL), the local node of iBOL, and the resulting data are made available through the Barcode of Life Data Systems (<http://www.boldsystems.org/>). The NTI projects will provide georeferenced records of non-native

species (submitted to the Norwegian Biodiversity Information Centre). In addition, the barcodes provided from these projects, is available in BOLD and can be used within activities on metabarcoding and eDNA. All collected material are stored and curated in the university museums' collections, for documentation and future research.

Examples of NTI projects providing species inventories and barcode databases: The project COPCLAD (2015–2017, IMR) performed an inventory of planktonic copepods and Cladocera in Norwegian waters. The project HYPCOP (2020–2022; IMR) is focusing on the diversity of copepods in the hyperbenthic marine habitats in Norwegian waters and will build a reference DNA-database on primarily harpacticoids. The project “Hardbunnsfauna” (2019-2022, Natural History Museum, UiB) will provide new knowledge on the distributions of marine invertebrate species in shallow hard bottom sites. Special focus is on sponges, bryozoans and ascidians. The project also contributes with DNA barcodes from a variety of species, which are submitted to the BOLD database.

Information on all marine projects within the Norwegian Taxonomy Initiative can be found at:

<https://www.artsdatabanken.no/Pages/195803/Hav>

2.14 - Summary of monitoring and mapping in Norwegian Waters

Monitoring of king crab and snow crab is mostly with the purpose of stock assessment for advice given to the management of the fisheries, but also to follow population development and dispersal in the Barents Sea and Norwegian Sea. Pink salmon is monitored through a network of activities along the entire coast. Pacific oyster has been monitored in its distribution area along the coast of Norway through a Nordic network and the project stops in 2021. Likewise, the pathogens in native and introduced oyster has been subject to monitoring. The red algae *Agarophyton vermiculophyllum*, which have caused problems in many countries, has been mapped along the coast of the Oslofjord south to Kristiansand. However, this mapping is very limited, as only one area has been mapped each year and the abundance of the species has only been recorded as presence/absence. The abundance of the species is only recorded properly at three sites. Alien species of zooplankton is occasionally detected in general monitoring programs of zooplankton. Additionally, is alien jelly fish reported in citizen science projects.

The Norwegian coast is the second longest in the world and mapping and monitoring of introduced species is time consuming and costly. About ¼ of the coast have been mapped with the purpose of getting a general picture of distribution of established species and possible newcomers. Each area has only been mapped once in a 10-year time-period and most sites have not been revisited. Several door knocker species may already be established along the coast, but the low mapping and monitoring effort have not been able to conclude on this. We have also little knowledge on the distribution on recent newcomers like *Hemigrapsus takanoi*, *Grateloupia turuturu* and *Didemnum vexillum* as no determined mapping program is yet in place. The round goby *Neogobius melanostomus* is present both on the Swedish west coast close to the Norwegian border and in Denmark and is expected to arrive here soon or is already present without being detected yet. In summer 2021 a small pilot study was performed on several sites in the Oslofjord, but this activity might not be enough to detect the species potential presence in Norwegian waters.

New species may be detected through general biodiversity projects, and by increasing our knowledge on general biodiversity we also increase our knowledge on alien species.

We also have little knowledge on the species associated with specific vector into the Norwegian coast such as transport water for cleaning fish, floating debris and species on ships. Biofouling communities on vessels

arriving from foreign ports have only been performed in a handful of vessels.

3 - Vectors for introduction of NIMS into the Norwegian coast

3.1 - Ballast water

Following more than a decade of work at the MEPC (Marine Environmental Protection Committee, IMO), the Ballast Water Management Convention was adopted in 2004. There were, however, numerous unsettled issues to be finalized in the years to come. The convention should enter into force 12 months after ratification by a minimum of 30 States, representing 35% of world merchant shipping tonnage. This requirement was met when Finland ratified the convention on 8th September 2016, hence the convention entered into force in 2017. By this time, most of the practical and regulatory remaining issues in the original convention had been resolved. The Convention requires all ships to implement a ballast water management plan. All ships must carry a ballast water record book and are required to carry out ballast water management procedures to a given standard. Parties to the Convention are given the option to take additional measures which are subject to criteria set out in the Convention and to IMO guidelines.

Ballast water treatment was scheduled in a two-tier approach. While allowing for a substantial “grace-period” (The global shipyard capacity simply was too small to facilitate retrofitting in existing vessels fast enough), an alternative standard (D1) was adopted. See infographics, Figure 19. The D1-standard requires the vessel to exchange ballast water at a distance > 200 nautical miles offshore, and preferably at depths > 200m. The requirements are specified in “Guidelines G6” in the Convention (2017 Guidelines for ballast water exchange (G6) (resolution MEPC.288(71)).




Figure 19. Infographics of the implementation of the Ballast Water Management Convention (Source: IMO).

New vessels built after 8th September 2017 must meet the D2 standard.

The D-2 standard specifies that ships can only discharge ballast water that meets the following criteria:

- less than 10 viable organisms per cubic metre which are greater than or equal to 50 micrometres in minimum dimension
- less than 10 viable organisms per millilitre which are between 10 micrometres and 50 micrometres in minimum dimension
- less than 1 colony-forming unit (cfu) per 100 millilitres of Toxicogenic *Vibrio cholerae*
- less than 250 cfu per 100 millilitres of *Escherichia coli*
- less than 100 cfu per 100 millilitres of Intestinal Enterococci

Older vessels must also meet the D2 when they are having

- IOPPC Renewal survey after 8th September 2019 A ship undergoing a renewal survey linked to the ship's International Oil Pollution Prevention Certificate after 8th September 2019 will need to meet the D2 standard by the date of this renewal survey.
- IOPPC Renewal survey between 8 September 2017 and 8 September 2019. - If the previous IOPPC renewal survey was between 8 September 2014 and 8 September 2017, then the ship must comply with D2 standard by this renewal survey. If the previous IOPPC renewal survey was before 8 September 2014, then the ship can wait until the next

renewal survey (which will be after 8 September 2019).

While vessels built after 8th September 2017 are required to meet the D2 standard, older vessels (not having to retrofit in conjunction with IOPPC renewal surveys (see above) may still operate under the D1 standard until the end of the “grace period” (8th September 2024). Areas for ballast water exchange (Figure 20) in compliance with the D1 standard are specified in the Norwegian by-law: “Forskrift om ballastvannbehandling på skip og flyttbare innretninger.” (Anon 2015).




Figure 20. Areas for ballast water exchange along the Norwegian coast (Map from Norwegian Maritime Authority).

While it is anticipated that the Ballast Water Management Convention will reduce the risk for NIS introductions, there are several types of vessels that are exempt from the Convention.

In addition to commercial vessels < 400 gross tons, cf. Regulation E-1.1., war ships, naval auxiliary, or other ships owned and operated by a state and used only on government non-commercial service, as stated in Article 3.2(d) of the convention. While the volume of ballast water carried by the bulk of these vessels are modest, the high number, and the unrestricted sailing pattern still represent risk for further spread of NIS.

3.2 - Biofouling on vessels

In 2011 guidelines for the control and management of ships' s biofouling to minimize the transfer of invasive aquatic species were presented by IMO (MEPC 62/24-2011). The guidelines include actions like biofouling management plan and record book, anti-fouling system installations and maintenance, and in-water inspection, cleaning and maintenance. The idea was to test out these guidelines on a voluntary basis before mandatory regulations were implemented. Currently there is ongoing work in IMO to assess these experiences and develop standardized regulations.

With the ballast water under better control, biofouling on ships probably is the vector with highest risk of bringing new species to Norwegian waters. From being a country with mostly fishing boats and smaller cargo up to the 1970's the development of the oil and aquaculture industry, the increased demand for imported goods and increased tourist activity has led to a dramatic change in the vessel intensity along the Norwegian coast the last five decades.

3.3 - Analyses of shipping activity in Norwegian ports

Voyage origin (last port call) were assembled from port history in Marine Traffic for vessels (n = 157 641) entering 143 Norwegian ports in the period October 2020 to November 2021. Port call in these categories were analysed: Cargo, tankers, tugs and special crafts (including all types of vessels related to the oil industry), fishing vessels and leisure crafts. Passenger vessels were excluded from the study, since most passenger vessels in this period were of national origin due to Covid-19 restrictions. Data for Ro-ro passenger vessels to ports that have ferries to Sweden and Denmark were assembled. To get a picture of the cruise traffic into a port in a normal year, data for port calls in 2019 were provided from Bergen Port Authorities, as Bergen port is one of the busiest cruise ports in Norway. Norwegian ports that received > 500 arrivals per year were included in the study. Total number of arrivals and arrivals from foreign ports for each vessel category were counted. Origin port for all vessels from foreign ports were identified to country for ports that received > 5 foreign arrivals per year. Marine traffic only gives historical data for port calls back one year and yields a maximum of 500 vessels in each category. For busy ports the number of 500 arrivals were exceeded in a variable time span for most vessel categories. The number of days in this timespan were noted and a mean daily rate of arrivals in this period were calculated and used to estimate a total number of annual arrivals in each category. Number of foreign arrivals were calculated in the same way for ports with > 500 arrivals. Data on shipping intensity is retrieved from European Marine Observation and Data network (EMODnet: <https://emodnet.ec.europa.eu/en>).

3.3.1 - Cargo

The group cargo includes vessels from small carriers to large container ships. A total estimated number of 65 500 cargo vessels arrived annually in Norwegian ports in 2020-2021. The highest total number of cargo arrivals was in Haugesund (5703), followed by Tromsø (2724), Florø (2500), Måløy (2607) and Ålesund (2253). Oslo port had the highest number of foreign cargo arrivals (925), followed by Porsgrunn (550), Haugesund (547) and Tananger (530) (Figure 21). (Detailed data is available in Appendix 1, Table 1).




Figure 21. (Left): Annual number of arrivals of cargo vessels into Norwegian ports. Shipping intensity for this group shown as routes/km²/year. (Right): Annual number of foreign cargo arrivals into Norwegian ports. Annual arrivals are estimated for ports which receives >500 arrivals a year (Data from Marine Traffic and EMODnet).

3.3.2 - Tankers

The group tankers include vessels from small harbour tankers to large oil and LNG tankers. A total estimated number of 17 200 tankers arrived annually in Norwegian ports in 2020-2021. The highest estimated number of tanker arrivals was in Mongstad port (2607) of which 1642 of the arrivals were from a foreign port followed by Rafnes (542) of which 434 was from a foreign port and Kårstø (500) of which 467 where of foreign origin. Several other ports had a high number of foreign tankers such as Fredrikstad (206), Slagen (196), Sture (183), Porsgrunn (180) and Elnesvågen (150) (Figure 22). (Detailed data is available in Appendix 1, Table 3).




Figure 22. (Left): Annual number of arrivals of tankers into Norwegian ports. Shipping intensity for this group shown as routes/ km²/year. (Right): Annual number of foreign tanker arrivals into Norwegian ports. Annual arrivals are estimated for ports which receives >500 arrivals a year (Data from Marine Traffic and EMODnet).

3.3.3 - Tugs and special crafts

The group 'tugs and special crafts' contains various types of vessels; small harbour tug, rescue boats, large international tugs and crane vessels, barges, all types of support vessels for the oil industry. A total estimated number of 144 848 'tugs and species craft' port calls annually in Norwegian ports in 2020-2021. As the map of route density shows most of the long-distance traffic is going between Norwegian ports and the oil fields in the North Sea and the Norwegian Sea (Figure 23). The highest estimated number of arrivals in this group was in Oslo (22813), followed by Brevik (14038) and Tromsø (4803). Oslo port had the highest number of foreign arrivals (1506), followed by Bergen (155), Brevik (112), Tananger (109) and Stavanger (102) (Figure 23). (Detailed data is available in Appendix 1, Table 2).




Figure 23. (Left): Estimated annual number of arrivals of tugs and special crafts into Norwegian ports. Shipping intensity for this group shown as routes/ km²/year. (Right): Percent annual foreign tugs and special crafts arrivals into Norwegian ports. (Data from Marine Traffic and EMODnet).

3.3.4 - Fishing vessels

The group fishing vessels contains small local fishing boats that goes in and out of the same harbour frequently to larger trawler. A total estimated number of 104 926 fishing vessels arrived annually in Norwegian ports in 2020-2021, of which 1,3 % was of foreign origin. As the map of route density shows there is high activity along the entire coast of the North Sea and the Norwegian Sea (Figure 24).

The highest estimated number of arrivals of fishing vessels in this group was in Tromsø (10139) of which none was of foreign origin. The same is the case for larger fishing port ranging from 3967 to 2340 annual arrivals such as Båtsfjord, Vardø, Vadsø, Honningsvåg, Rørvik, Måløy, Kristiansund and Ålesund (Foreign share 0-1%).

The highest share of foreign fishing vessels was in Kirkenes in the north (30%) where most of the boats came from ports in Northern Russia and a few from Iceland and Lithuania. Kristiansand in the south have 10 % share of foreign fishing boats most of them coming in from Denmark. Ports in the Skagerrak and North Sea area receives a relatively low share of fishing vessels, most of them coming from the other Nordic countries but also GB, Ireland, Iceland, Faroe Islands and Germany (Figure 24).




Figure 24. Estimated annual number of arrivals of fishing vessels into Norwegian ports. Shipping intensity for this group shown as routes/ km²/year. (Data from Marine Traffic and EMODnet).

3.3.5 - Passenger vessels and leisure crafts

3.3.5.1 - Ferries

There is frequently ferries (ro-ro passenger) going back and forth to Denmark, Sweden and Germany and into Norwegian ports. The ports of Tananger, Kristiansand and Larvik had respectively 212 and 500 arrivals from Hirtshals in Denmark. Oslo received 51 ferries from Fredrikshavn in Denmark and 168 from Kiel in Germany. The port of Sandefjord received 498 arrivals from Strømstad in Sweden.

3.3.5.2 - Leisure craft

The majority of port calls in this group are boats going in and out of the marina daily or travellers along the coast during summer season. The ports along in the Skagerrak area and Oslofjord received also some travellers from Sweden and Denmark. Occasionally some port received boats from Iceland, Faroe Island, GB, Ireland, The Netherlands, Germany and the Baltic Sea. In a normal year with no Covid-19 restriction there is probably more leisure craft arrivals from abroad.

3.3.5.3 - Cruise ships

The cruise traffic in Norway has increased steadily the last decades from 1033 arrivals in 1993 to 2159 in 2018 (Dybedal 2018). The number of cruise passenger has in the same period been four-doubled, which means that the ships arriving is larger than before. Bergen is the busiest cruise port in Norway, followed by Geiranger,

Stavanger, Ålesund, Flåm, Tromsø, Nordkapp and Oslo.

The last two years the activity has been low in all ports due to Covid-19 restrictions. Many of the cruise ships are sailing in warmer parts of the world during winter and returns to European port to get ready for cruises up north in spring/early summer.

To get a picture of the origin of cruise ships to the Norwegian coast an analysis of the last port before arrival for 378 cruise ships into Bergen in 2019 is performed. Most of the cruise ships came from other Norwegian cruise harbours, which reflects that passengers are flown in and the ships goes back and forth along the coast (Figure 25). The foreign arrivals were from Germany, Denmark, Belgium, the Netherlands, Great Britain, Iceland, and Faroe Islands. 12 ships came from the northeast coast of the US (not shown in map in Figure 25).



Figure 25. Last port call before arrival in Bergen for 378 cruise ships in 2019 (Data and map: Bergen Port Authorities).

3.3.6 - Svalbard

The two ports in Svalbard, Longyearbyen og Ny-Ålesund, had 2117 annual port calls in 2020-2021 (Figure 26). Seven of the vessels came from ports outside Norwegian territory, while 113 came from mainland Norway. The seven foreign vessels were leisure crafts from Iceland, Belgium and Poland. As figure 26 shows there is high fishing activity around Svalbard, but few boats go into the ports.




Figure 26. Estimated annual number of arrivals of cargo (top left), tankers (top right), Special ships (bottom left) and fishing vessels (bottom right) into Svalbard ports in 2020-2021. Shipping intensity for is shown as routes/ km 2/year. (Data from Marine Traffic and EMODnet).

3.3.7 - The ports behind the ports

The Norwegian coast is long and for smaller vessels coming from abroad it is more and more likely that they have stopped in several Norwegian harbours before entering northern Norway. This will of course be reflected as bias in the data and probably underestimate the vector as going northwards along the coast. Larger boat like tankers and large cargo vessels with a specific task in one harbour is more likely to go directly from international waters into Norwegian ports. Furthermore, many vessels come from large ports in UK and western Europa such as Aberdeen, Rotterdam and Amsterdam. Though many of them most likely have had an operational area the last year which covers a much larger geographical span. To uncover the full picture, one must assemble historical data on each vessel in Marine Traffic, which is an almost impossible task when the number of arrivals exceeds 150 000. To elucidate this problem, we have picked 20 of the last arrivals from Rotterdam into Rogaland County in the groups cargo and tankers which clearly shows that the geographical origin of species on arriving vessels might be much larger (Figure 27). The number of vessels operating in area exceeds 40 since most ships have been in several areas. An analysis of the ports visited during 2019 by cruise ships that visited Bergen port that year also shows the extent of wider operation area of vessels. The cruise ships have visited ports all around the world during that year before entering European waters. Each vessel had a mean number of 75 port call during 2019 (Figure 28).




Figure 27. Operational area the last year of 40 randomly selected vessels (cargo & tankers) entering ports (Haugesund, Tananger and Kårstø) in Rogaland from Rotterdam in 2021 (Data from Marine Traffic).




Figure 28. Ports visited during the year by 378 cruise ships that entered Bergen in 2019 (Data from <http://crew-center.com/cruise-ship-ports-schedules>).

3.3.8 - Geographical distribution and origin of foreign arrivals in Norwegian Counties

Number of annual arrivals is based on manually counted arrivals in each vessel group but gives a skewed picture since only the last 500 port calls are shown in Marine Traffic history. Given the rate of vessels coming from abroad are approximately the same during the year, an estimate of the number of foreign arrivals into ports with >500 arrivals a year is calculated (Figure 29).

Viken & Oslo is the county with the highest annual number of foreign arrivals (3738) in the groups cargo, tankers and special crafts, followed by Rogaland (3585) , Vestland (3197) and Vestfold & Telemark (2293). Svalbard had no records of vessels in this group coming from countries outside Norway, but 113 arrivals from mainland Norway. Rogaland had the highest number of cargo vessels coming from abroad, Vestland the highest number of tankers, while Viken & Oslo had the highest number of foreign special crafts (Figure 29).




Figure 29. Annual foreign arrivals (estimated for ports with >500 annual arrivals) in the groups cargo, tanker and special crafts in Norwegian Counties (Data from Marine traffic).

3.3.8.1 - Viken & Oslo, Vestfold & Telemark

These are the four counties surrounding the inner and outer Oslofjord. Viken and Oslo receives annually 3738 foreign arrivals in the groups cargo, tankers, tugs and special crafts. Cargo vessels represents the largest group with 1859 arrivals, followed by tugs and special craft (1536) and tankers (343). Vestfold & Telemark receives annually 2293 foreign arrivals in these groups. Cargo vessels represents the largest group with 1316 arrivals, followed by tankers (915) and tugs & special craft (62). For detailed information on each port and vessel type see Appendix 1, Table 1, 2, 3. For detailed data on origin of foreign vessels see Appendix 2, Table 1.

Cargo into this region mainly originates from European countries like the Netherlands, Belgium, Germany, Spain, UK, Sweden and Denmark and countries in the Baltic area. The industrial ports of Brevik and Porsgrunn receives some cargo from the Middle east, North Africa, and Asia (Figure 30). Fredrikstad had some arrivals of cargo from Brazil and Canada (Figure 30).

Tankers into this area mainly originates from European countries like the Netherlands, Belgium, Germany, Spain, UK, Sweden and Denmark and countries in the Baltic area, as well as a fair share of arrivals from USA, India, Oceania, and Brazil (Figure 31).

Tugs and special crafts of foreign origin comes mainly into the port of Oslo from Sweden and Denmark, the Netherlands, Belgium, Germany and Poland (Figure 32). The port of Slagen receives frequently vessels from Denmark and the port of Drammen from the Netherlands, Germany, and Sweden.




Figure 30. Origin (last port) of 3175 cargo vessels arriving in ports in the counties Oslo, Viken, Vestfold & Telemark in the period October 2020-2021 (Data from Marine Traffic).




Figure 31. Origin (last port) of 1258 tankers arriving in ports in the counties Oslo, Viken, Vestfold & Telemark in the period October 2020-2021 (Data from Marine Traffic).




Figure 32. Origin (last port) of 1598 tugs and special crafts arriving in ports in the counties Oslo, Viken, Vestfold & Telemark in the period October 2020-2021 (Data from Marine Traffic).

3.3.8.2 - Agder

The ports in Agder have a relatively low annual share of foreign arrivals in the groups cargo (518), tankers (8), and tugs and special crafts (62). For detailed information on each port and vessel type see Appendix 1, Table 1, 2, 3.

Cargo into this region mainly originates from European countries like the Netherlands, Belgium, France, Spain, Germany, UK, Sweden and Denmark and countries in the Baltic area. Kristiansand port receives several cargo vessels from Canada. Eydehavn and Grimstad additionally have some arrivals from the Mediterranean and North African countries (Figure 33).

Farsund, Mandal, Kristiansand and Arendal are the port that receives most foreign tugs and special craft, mostly arriving from Denmark, Germany, Sweden, Finland, and UK (not shown in map).

The only port which received tankers from abroad were Farsund, with eight arrivals from Sweden (not shown in map).




Figure 33. Origin (last port) of 487 cargo vessels arriving in ports in Agder County in the period October 2020-2021 (Data from Marine Traffic).

3.3.8.3 - Rogaland

Rogaland receives annually 3738 foreign arrivals in the groups cargo, tankers, tugs and special crafts. Cargo represents the largest group with 2402 arrivals, followed by tankers (749), and tugs and special crafts (434). For detailed information on each port and vessel type see Appendix 1, Table 1, 2, 3. For detailed data on origin of foreign vessels see Appendix 2, Table 1.

The majority of cargo vessel comes in from the Netherlands, followed by Germany, UK, Sweden, Denmark and the Baltic Sea. Some vessels come from further south in Europe, France, Spain and the Mediterranean, also on the African side and South Africa (Figure 34).

The oil terminal at Kårstø is the port with the highest traffic of tankers with annual arrivals of 500 vessels, of which 91 % comes from a port abroad. Tankers into this port arrives mainly from the Netherlands, France, Germany, UK (including Ireland), Sweden and the Baltic Sea. Some vessels also arrive from the Mediterranean, Africa, North and South America (Figure 35).

The ports in Haugesund and Husøy receives most special crafts from UK and Denmark with some vessels coming from the Netherlands and Germany, USA, Angola and South Africa. The port of Stavanger and the nearby ports receives most special crafts from UK and the Netherlands, with some arrivals from other European, Nordic and Baltic countries, the Mediterranean, Africa, India and USA (Figure 36).




Figure 34. Origin (last port) of 2402 cargo vessels arriving in ports in Rogaland County in the period October 2020-2021 (Data from Marine Traffic).




Figure 35. Origin (last port) of 4342 tankers arriving in ports in Rogaland County in the period October 2020-2021 (Data from Marine Traffic).



Figure 36. Origin (last port) of 434 tugs and special crafts arriving in ports in Rogaland County in the period October 2020-2021 (Data from Marine Traffic).

3.3.8.4 - Vestland

Vestland receives annually 3197 foreign arrivals in the groups cargo, tankers, tugs and special crafts. Tankers represents the largest group with 1913 arrivals, followed by cargo (946), and tugs and special crafts (338). For detailed information on each port and vessel type see Appendix 1, Table 1, 2, 3. For detailed data on origin of foreign vessels see Appendix 2, Table 1.

Florø is the port that receives most foreign cargo vessels, followed by Odda, Måløy and Sløvåg. Cargo vessel into this county mainly arrives from the Netherlands, followed by UK, Germany, Sweden, France, Denmark and the Baltic Sea. Odda has a fair share of vessels from Spain, Portugal, and North Africa. The industrial ports of Høyanger and Husnes have several arrivals from Brazil (Figure 37).

The port of Mongstad has the highest traffic of tankers with annual arrivals of 2607 vessels, of which 63 % comes from a port abroad. The oil terminal at Sture is the second largest port in this group with 203 arrivals where 90 % comes from abroad. Tankers into these ports arrives mainly from the Netherlands, France, Germany, UK (including Ireland), Sweden, Denmark and the Baltic Sea. Some vessels also arrive from the Mediterranean, Africa, North & South America, the Arabic Gulf and Asia (Figure 38).

Bergen is the port with most arrivals of foreign tugs and special crafts, followed by Florø, Mongstad, Måløy and Ågotnes. Most of the vessels arrives from UK and Denmark with some vessels coming from the Netherlands

and Germany, Sweden, Denmark and the Baltic Sea. Note that the records of vessels from Africa in the port of Torangsvåg is based on one recorded vessel from Angola and is mistakenly included in the data (Figure 39).




Figure 37. Origin (last port) of 946 cargo vessels arriving in ports in Vestland County in the period October 2020-2021 (Data from Marine Traffic).




Figure 38. Origin (last port) of 1913 tankers arriving in ports in Vestland County in the period October 2020-2021 (Data from Marine Traffic).




Figure 39. Origin (last port) of 338 tugs and special crafts arriving in ports in Vestland County in the period October 2020-2021. Note that the records of vessels from Africa in the port of Torangsvåg is based on one recorded vessel from Angola and is mistakenly included in the data (Data from Marine Traffic).

3.3.8.5 - Møre & Romsdal

Møre & Romsdal have relatively few arrivals from foreign ports (508), but this far north on the coast there is a high probability that vessels arriving from a Norwegian port have a wider international operating area. The area receives 270 cargo vessels, 194 tankers and 44 tugs and special crafts. For detailed information on each port and vessel type see Appendix 1, Table 1, 2, 3. For detailed data on origin of foreign vessels see Appendix 2, Table 1.

Ålesund is the port which receives most cargo vessels (2253) of which 4 % is from a foreign port followed by Kristiansund (1534) with only 1% of foreign origin. The port at Averøya receives most foreign cargo vessels (92) followed by Ålesund (90) and Sunndalsøra (37). The ports in this area mainly receives vessels from western Europe, Sweden and Denmark, the Baltic Sea. Ålesund had some arrivals from Egypt and north Africa. Sunndalsøra had 19 arrivals from Brazil and 2 from USA (Figure 40).

The port of Elnesvågen has the highest number of foreign tankers arriving (150) mainly from the Netherlands, Germany, and Sweden but also some from other European ports, the Baltic and North Africa. Kristiansund and Averøya have some arrivals of tankers from European, Nordic and Baltic ports (Figure 41).

The only two ports that had arrivals of foreign tugs and special crafts was Ulsteinvik with seven vessels and Ålesund with 15 vessels from UK, Germany, Denmark, Poland and Turkey (not shown in map).




Figure 40. Origin (last port) of 270 cargo vessels arriving in ports in Møre & Romsdal County in the period October 2020-2021 (Data from Marine Traffic).




Figure 41. Origin (last port) of 194 tankers arriving in ports in Møre & Romsdal County in the period October 2020-2021 (Data from Marine Traffic).

3.3.8.6 - Trøndelag

Trøndelag had relatively few arrivals from foreign ports (228), but this far north on the coast there is a high probability that vessels arriving from a Norwegian port have a wider international operating area. The area receives 162 cargo vessels, 45 tankers and 21 tugs and special crafts. For detailed information on each port and vessel type see Appendix 1, Table 1, 2, 3. For detailed data on origin of foreign vessels see Appendix 2, Table 1.

Cargo vessels from foreign ports in this county is only coming into the ports of Orkanger (71), Trondheim (35) and Verdal (56). Origin of cargo vessels into these ports are mainly the Netherlands, Spain (Verdal), UK, Sweden, Denmark and Poland (Figure 42).

Trondheim is the only port which received >5 foreign tankers in this county (41), most of them arriving from Sweden and some from the Netherlands (not shown in map).

Verdal is the only port which received >5 foreign tugs and special craft in this county (12), most of them arriving from the Netherlands, Germany, Denmark and Spain (not shown in map).




Figure 42. Origin (last port) of 162 cargo vessels arriving in ports in Trøndelag County in the period October 2020-2021 (Data from Marine Traffic).

3.3.8.7 - Nordland

Nordland have relatively few arrivals from foreign ports (806), but this far north on the coast there is a high probability that vessels arriving from a Norwegian port have a wider international operating area. The area receives 738 cargo vessels, 52 tankers and 16 tugs and special crafts. For detailed information on each port and vessel type see Appendix 1, Table 1, 2, 3. For detailed data on origin of foreign vessels see Appendix 2, Table 1.

Cargo vessels in this county mainly comes from countries around the North Sea, Nordic and Baltic countries. The industrial port in Glomfjord additionally receives vessels from northern Russia and Morocco. Mosjøen have also several cargo vessels coming in from Spain, Turkey, USA and Brazil. The port of Narvik, which have been shipping out minerals from Kiruna in more than 100 years, received additional vessels from Turkey, Morocco, Saudi Arabia, USA, India and Egypt. Sortland and Stokkmarknes had a low number of visits but a fair share of cargo vessels from northern Russia (Figure 43).

Mosjøen, Bodø and Glomfjord are the only ports which received >5 foreign tankers in this county, most of them arriving from Sweden and Denmark and some from other European and Baltic countries. Glomfjord had

additional one port call from Trinidad (Figure 44).

None of the 21 ports examined in Nordland had >5 foreign tugs and special craft, and the origin of the 16 vessels in this group is therefore not noted.




Figure 43. Origin (last port) of 738 cargo vessels arriving in ports in Nordland County in the period October 2020-2021 (Data from Marine Traffic).




Figure 44. Origin (last port) of 52 tankers arriving in ports in Nordland County in the period October 2020-2021 (Data from Marine Traffic).

3.3.8.8 - Troms & Finnmark

Troms & Finnmark had the lowest number of arrivals from foreign ports (126), but this far north on the coast there is a high probability that vessels arriving from a Norwegian port have a wider international operating area. The area receives 126 cargo vessels, 30 tankers and 32 tugs and special crafts. For detailed information on each port and vessel type see Appendix 1, Table 1, 2, 3. For detailed data on origin of foreign vessels see Appendix 2, Table 1.

Cargo vessels from abroad mainly arrived in Kirkenes (41), Båtsfjord (45), Tromsø (22) and Bergsneset (11). The vessels mainly came from northern Russia and the Netherlands, Iceland and the Faroe Islands and from Denmark and the Baltic Sea (Figure 45).

Tromsø and Harstad are the only ports that received >5 tankers from abroad, mainly originating from Sweden, but also some port calls from the Netherlands UK and Denmark (not shown in map).

Kirkenes is the only port that received >5 tugs and special craft from abroad, mainly originating from northern Russia and the Netherlands (not shown in map).




Figure 45. Origin (last port) of 126 cargo vessels arriving in ports in Troms & Finnmark County in the period October 2020-2021 (Data from Marine Traffic).

3.4 - Import and sale of living seafood

Handling of living imported seafood is regulated by law (Anon. 2015). Keeping imported organisms in containers requires measurements to avoid escape and emission of larvae in the waste water. Moreover is the seller responsible to inform customers that this is an alien species and how to prevent it from entering the environment.

Live Pacific oysters (*Crassostrea gigas*) are imported from France and The Netherlands to supply the Norwegian market for live oysters. There is no control of oyster diseases, as the oysters are classified as food and destined for “the table”, without repacking or re-laying. The oysters come from dispatch centres and are usually packed in units that fits the restaurant or fresh seafood market. This minimizes the risk associated with re-laying in tank units without proper water treatment or in open waters. It is difficult to get hold of information on which species of bivalves are imported for consume to the Norwegian market. Well assorted seafood shops offer living oysters, razor clams, cockles and various hard shell clams etc. but it is difficult to know whether they are of Norwegian origin without a more throughout investigation (Figure 46).

The extensive import of living American lobster into Europe, has led to several records of this species in Norway. Lobsters were often found with stripped claws and had most likely escaped from containers in the sea. In 2016 it became illegal to import living oysters to Norway. One company has a dispensation from this

regulation, but the living lobster is kept in escape secured containers and waste waters is filtered and UV-radiated. The lobster is cooked before it goes to the market. After 2017 there has been no records of American lobsters in Norway, but wild populations might be established in areas with many records or where hybrid eggs have been spawned. We have no knowledge about other crustaceans imported live to Norway.



Figure 46. Pacific oysters, razor clams and European flat oyster in a seafood shop (Photo: Fjellskål, Bergen).

3.5 - Transport of live cleaner fish

Parasitic salmon lice represent a major problem in Norwegian aquaculture and cause economic losses in the aquaculture of Atlantic salmon and rainbow trout. It also has a significant impact on wild populations of salmonids, particularly sea trout in coastal areas with aquaculture activity. Several pharmaceuticals have been used for treatment of salmon lice infestations, but over time the lice have developed resistance to these treatments, and there is a growing concern regarding the environmental impact of chemical and pharmaceutical treatments.

There is a strong incentive to find alternative methods of de-lousing. The use of cleaner fish, which pick lice

from the skin of salmon and rainbow trout, has become an important tool (although this is associated with other problems). Lumpfish and several species of wrasses are transferred to the net pens and used as cleaner fish. To supply the fish farms with cleaner fish, extensive fisheries for wrasses have developed along the Swedish west coast and in southern and western Norway. In addition, there is aquaculture of ballan wrasse and lumpfish in Norway.

The Norwegian fish farming industry uses around 60 million cleaner fish annually. Most wrasses used in the fish farms are caught locally, but around 25 % are moved from South-eastern Norway and Southwestern Sweden and translocated to fish farms further north. Farms in mid-Norway are the main “importers” of live wrasses. In 2020 around 18 million wrasses were fished in Norway. In addition, around one million were fished in Sweden. When cleaner fish are introduced and escape outside their natural distribution range in Norway they are defined as regionally alien fish, which is the case for corkwing wrasse *Symphodus melops* and rock cook *Centrolabrus exoletus* (Norwegian Biodiversity Information Centre 2018).

3.5.1 - Risk factors related to translocation of live cleaner fish

Translocation of cleaner fish between distant populations can affect native populations through disease transfer and result in irreversible genetic changes if they escape. The risk associated with import and translocation of fish has been assessed by The Norwegian Scientific Committee for Food and Environment (Vitenskapskomiteen for mat og miljø, VKM) (VKM 2017, 2019), and by the Institute of Marine Research (Grefsrud et al. 2021a,b). Risk is associated both with the transfer of the fish and the transport water.

To avoid transmission of disease, it is as a general principle not possible to mix wild and farmed animals, within a farming unit, or even keep wild caught and farmed fish in proximity. Cleaner fish and salmonids may share pathogens (Erkinharju et al. 2020). Pathogens may change virulence, and polyculture with a mixture of susceptible hosts and new pathogens may trigger host-switching and adaptation to new species and environment. Repetitive long-distance transport of wild caught wrasses with unknown health status represents a risk of disease transfer. The current farm practice with its open design increases the risk of transmission between farmed and wild fish. In addition, the high density of hosts in the fish farms may also increase disease prevalence.

Reports from the Institute of Marine Research have focused on pathogenic viruses, and particularly emphasized the risk of introducing viral haemorrhagic septicaemia virus (VHSV). It has also been recommended by VKM in 2019 to start screening of cleaner fish for VHSV (VKM 2019). VHSV adapts to new hosts and habitats. VHSV is a notifiable disease that is present in wild fish (http://web.oie.int/eng/maladies/en_classification2010.htm). Outbreaks of VHS have been reported in wrasses in the Shetland Isles (Munro et al. 2015) and lumpfish in Iceland (Guðmundsdóttir et al. 2019). It is known that VHSV can transmit via food (Ahne 1980, Schönherz et al. 2012). Small wrasses are sometimes eaten by salmon or rainbow trout, particularly during the period of starvation before slaughter. Salmonids may thus be exposed to VHSV through infected wrasse. Another example is the Nerval necrosis virus (NNV) has been detected in wild Ballan, goldsinny and corkwing wrasse (Korsnes et al. 2017). NNV has been described from several marine fish species and is known from disease outbreaks in farmed Atlantic cod and Atlantic halibut (Patel et al. 2007; Grotmol et al. 1997). Different variants of the virus have been found in southern and northern parts of the distribution range of the wrasses, and translocations may lead to an import of the southern virus variant to northern areas.

Recent genetic studies show that cleaner fish escape and reproduce with local populations in the northern range, which can result in genetic changes and potentially reduced fitness in the local populations. Three species are translocated; the ballan, corkwing and goldsinny wrasses. These species live in shallow water

habitats and in geographically distinct populations. Recent studies have revealed significant genetical differences between these populations. Seljestad et al. (2020) found a clear genetic break dividing Scandinavian populations of Ballan wrasse into north-western and south-eastern groups.

For corkwing wrasse, there is a large genetic break between Scandinavian and UK populations, and additionally a strong population structure within Scandinavia (Blanco Gonzalez et al. 2016, Robalo et al. 2012, Knutsen et al. 2013). The goldsinny wrasse populations are also clearly genetically divided across the North Sea (Jansson et al. 2017; 2020) but in contrast to the corkwing and ballan wrasse, goldsinny has a relatively weak population structure, but with a clear pattern of isolation-by-distance (Jansson et al. 2017).

Wrasse can escape from the salmon pens through holes in the net, small fish slipping through the mesh (Svåsand et al. 2017, Woll et al. 2013). Recent genetic studies show that cleaner fish in Trøndelag are escaping and reproducing with local populations (Jansson et al. 2017, Faust et al. 2018). Currently, it is permitted to import wrasse from Sweden during the spawning period – which probably increase the likelihood of genetic introgression.

No studies have been performed on the transport water. Repeated long-distance transport and release of untreated water from shallow water areas represents a hazard of moving the entire microbiota from the donor habitat to the recipient habitat. This includes plankton organisms as well as propagules from benthic algae and larvae of alien mollusc, crustacean and fish species.

3.6 - Trade of species for saltwater aquaria

There is no need for permission to import marine plants, invertebrates or fish to be kept in aquaria where they cannot escape, however, there is a need to inform the Norwegian Environmental Agency (Anon. 2015). Most saltwater species in aquarium shops in Norway come from tropical or subtropical regions and will therefore most likely not be able to establish in Norway, should they be released. Based on the low probability of warm-water marine organisms surviving under Norwegian conditions if released, the risk of new invasive marine species coming from aquarium trade is considered low. However, there can be exceptions, and the possibility of buying species on the web makes it difficult to get an overview of the magnitude of this, thus better data and reporting systems is needed to increase the knowledge and reduce the uncertainty on possible impact of this vector.

3.7 - Species transported by floating debris

Attention to the ability of alien species to hitch hike with floating debris was brought to attention when a mass event of non-native species crossing the Pacific Ocean with floating debris from the tsunami disaster in Japan in 2011 took place. A study counted 289 Japanese marine species on debris along the shore (Carlton et al. 2017). Taxa in the phyla Mollusca, Annelida, Cnidaria, Bryozoa, Crustacea, Porifera, Ascidia, Echinodermata and Pisces have survived for a six years journey across the ocean. Biota were attached or entangled in substances like boats, docks, buoys or beams.

Rafting of species on natural floats like kelp or fallen trees has been known from ancient times. However, floating debris from other human activity is an increasing problem in the oceans. Rech et al. (2016) consider floating debris to be of high importance for both dispersal of new marine species and for the secondary dispersal of established ones. The risk for establishment of NIMS on floating debris is dependent on the material and buoyancy of the debris, the donor region and on the conditions when arriving at the coast. On litter that has landed on the beach and dried out calcified and crustose organisms is more common. On litter still in the sea organisms from almost all phyla can be present (Figure 47).

In recent years several studies have investigated the biota on floating debris. (Gregory 2009, Campbell et al. 2017, Gündoğdu et al. 2017, Tutman et al. 2017). Kiessling et al. (2015) reports on 387 taxa, including pro- and eukaryotic micro-organisms, seaweeds and invertebrates found on floating litter.

A model-study of the origin of plastic litter, based on collected litter on seven beaches along the coast and at Svalbard, showed that most of the litter that comes ashore along the southern coast of Norway originates from the southern North Sea and Skagerrak area. The beaches at the western coast and northern coast comes from the Northern part of the North Sea and the Norwegians Sea. Litter arriving on the Svalbard beaches mostly originates from the areas around Svalbard and the Barents Sea (Strand et al. 2021).

In Norway no studies on species on marine debris has been conducted, but it can be of higher importance for primary and secondary spread of NIMS than anticipated.




Figure 47. Seaweeds, ascidians, and bryozoans growing on a buoy. Photo: Erling Svensen

3.8 - Transport of species with fishing gear

Transport of invasive species with fishing gear is well known for limnic species but less studied in the marine environment. Several adult species, larvae and algal propagules can probably survive in heaps of beams and

trawls when left in moist conditions. As there has been few studies on which species can survive such transport, we can only make a qualified guess on this matter. Several introduced seaweeds which reproduce clonally by fragments, such as *Dasysiphonia japonica* and *Agarophyton vermiculophyllum* (Nyberg & Wallentinus 2005, Husa et al. 2008) can probably survive in moist nets. Species of crustaceans, bryozoans, and ascidians as well. Of particular concern is the newly recorded species *Didemnum vexillum* which can probably survive on moist nets. Species like mussels, oysters, gastropods, sponges and crustaceans might also survive such conditions, while fish is probably less likely.

3.9 - Intentional introduction of species and import of species for aquaculture

Historically Norway has a very limited praxis of intentional introduction of species and import of species for aquaculture in the sea.

In 1988-1990 manila clam *Ruditapes philippinarum* was set out at six locations to study whether the species was a suitable species for aquaculture. Upon revisit of the sites in 1996 living animals were found, but there were no signs of recruitment (Mortensen & Strand 2000). However, species intentionally introduced to the marine environment in neighbouring countries have spread to the Norwegian coast.

Rainbow trout (*Oncorhynchus mykiss*), from North America, was introduced and released to lakes and rivers in Norway in the early 1900s to increase salmonid fisheries. Rainbow trout is also used in aquaculture in the coastal zone. Aquaculture of rainbow trout in Norway has increased since the 1990s and varies between 50 000 and 90 000 tonnes a year. Escapees of rainbow trout can interact with native salmonid fish by disturbing breeding areas and disperse parasites to the environment (Skaala et al. 2021).

In the early 1980's, permits were given to import Pacific oyster brood stock. The aim was to establish hatchery production of seed in two hatcheries and farming in oyster farms along the coast. The import was however continued, and relatively large amounts of imported oysters meant for human consumption were kept in farms or in dispatch centres without water treatments. Oysters were also re-laid outside one of the centres and have spawned (Wrange et al. 2010). Although the main invasion of Pacific oysters came with drifting larvae carried with the northward water currents, it is possible that re-laid oysters may have contributed to the establishment of feral populations. As oysters destined for human consumption are not subjected to health control, the oysters may carry oyster pathogens pose a disease risk for native bivalves.

New intentional introductions and import of aquaculture organisms to open systems in the sea are not likely to happen in Norway nowadays as this practice is regulated by law (Anon 2015). Permissions are given for aquaculture of alien species in escape secured land-based farms with treatment of waste water, such as Russian and Siberian sturgeon, scampi and abalone.

4 - Ecological impact assessment of vectors, species, and biogeography

In this chapter we take a closer look at pathways of introduction to Norway based on the information assembled in the Alien Species List of Norway (Norwegian Biodiversity Information Centre 2018). This list contains all alien species (within certain delimitations, see Sandvik et al. 2019) known to reproduce in the wild in Norway, in addition to selected door knocker species. The latter are species that do not currently reproduce (or even occur) in Norway but may be expected to do so in the near future (within 50 years). The Alien Species List does not only represent a catalogue of these species, but also lists the results of an ecological impact assessment, known and suspected pathways of introduction and spread, area of occupancy etc., for each species (Sandvik et al. 2020).

Pathways were recorded according to the subdivisions introduced by Hulme et al. (2008), adopted by the Convention on Biological Diversity (CBD 2014) and refined by Harrower et al. (2020). Regarding information on pathways in the Alien Species List, however, some pathway categories had been misinterpreted during impact assessments, as was evident from the fact that the pathway (sub)categories chosen were incompatible with the detailed descriptions given (Norwegian Biodiversity Information Centre 2018). For species for which sufficient information had been provided in the Alien Species List, the misinterpreted pathways have here been re-coded to the correct pathways following the detailed definitions by Harrower et al. (2020), see Box 1.

Box. 1 Re-coded pathways

The misinterpreted pathways were "other release" (*Eriocheir sinensis* was re-coded as "unaided dispersal"), "other escape" (*Ensis directus* was re-coded as "unaided dispersal"), "contaminant on animals" (*Crassostrea gigas* was re-coded as "escape from aquaculture"), "contaminant on plants" (*Dikerogammarus villosus* and *Watersipora subatra* were re-coded as "unaided dispersal"), "hitchhikers on boats" (*Oncorhynchus mykiss* was re-coded as "escape from aquaculture"; *Mnemiopsis leidyi* was re-coded as "ballast water"; *Amphibalanus amphitrite*, *Bonnemaisonia hamifera*, *Codium fragile*, *Bugula neritina*, *Cancer irroratus*, *Cryptonemia hibernica*, *Dasya sessilis*, *Schizoporella japonica*, *Tricellaria inopinata* and *Undaria pinnatifida* were re-coded as "hull fouling"; *Hemigrapsus sanguineus* and *Hemigrapsus takanoi* were re-coded as both "ballast water" and "hull fouling"; *Gracilaria vermiculophylla* was left unchanged because of insufficient information), "fishing equipment" (*Agardhiella subulata* was re-coded as "contaminant on animals"), "other stowaways" (*Aglaothamnion halliae* was re-coded as "escape from aquaria"; *Grateloupia turuturu*, *Rapana venosa* and *Ulva australis* were re-coded as "contaminant on animals"; *Agardhiella subulata* and *Bonnemaisonia hamifera* were re-coded as "hull fouling"; *Pachycordyle navis* and *Perophora japonica* were re-coded as "unaided dispersal"; *Crepidula fornicata* was re-coded as both "contaminant on animals" and "unaided dispersal"; *Colpomenia peregrina*, *Dasysiphonia japonica*, *Sargassum muticum* and *Tricellaria inopinata* were left unchanged because of insufficient information) and "artificial waterways" (*Cornigerius maeoticus* was re-coded as "unaided dispersal").

4.1 - Alien species currently reproducing in Norwegian waters

According to the Alien Species List, 37 alien marine species occurred in Norway by 2018, 17 of which were categorised as having a high or severe ecological impact (Table 1). Information on pathways of introduction was available for all species. A total of 80 pathways of introduction were reported, which amounts to an average of 2.2 pathways per species (range 1–4). All main pathways categories except corridors (interconnected waterways) were represented (Table 2). Stowaways and unaided spread were the dominant pathways of introduction. Pathways did not differ much between different taxonomic groups or between species with

high/severe impact and the remaining species (Table 2).

As regards pathways of secondary spread within Norway (i.e. following introduction), 44 such pathways were reported for 33 of the species, amounting to an average of 1.2 pathways per species (range 0–4). Unaided spread was the dominant pathway of secondary spread. Again, there were no pronounced differences between taxonomic groups or impact categories (Table 3).

Subcategories of pathways are summarised in Table 4. The most important subcategories of stowaways were ballast water, hull fouling and fishing equipment. This is in accordance with the fact that shipping is the most important factor for introduction of marine alien species to European seas (Nunes et al. 2014). A detailed summary for the species in the highest impact category ("SE") is provided in Table 5.

The geographical distribution of alien marine species shows that Southern and Western Norway is most exposed to introductions (Figures 49 and 50). The distribution is shown for all 39 alien species with individuals that are capable of "surviving in the wild" (categories C1–E sensu Blackburn et al. 2011; Figure 49) and the subset of 27 species "with individuals dispersing, surviving and reproducing at multiple sites" (only category E sensu Blackburn et al. 2011; Figure 50; data from Sandvik et al. 2019).

The temporal pattern of introductions of marine species to Norwegian waters does not display any saturation (Figure 51). To the contrary, decadal introduction rates of novel marine alien species have clearly been increasing from 1.1 ± 0.1 (1900–1953, N = 8) via 3.5 ± 0.1 (1953–1995, N = 15) to 7.5 ± 0.1 (1995–2014, N = 17) species per decade.

Table 1 . Ecological impact assessments of marine alien species known to reproduce in Norway by 2018. Tabulated by taxon and ecological impact category (NK = "no known impact", LO = "low impact", PH = "potentially high impact", HI = "high impact", SE = "severe impact"). (Source: Norwegian Biodiversity Information Centre , 2018).

Taxon	N	NK	LO	PH	HI	SE
Algae	10	-	2	3	-	5
- Bonnemaisoniales	1	-	-	-	-	1
- Ceramiales	5	-	2	2	-	1
- Gracilariales	1	-				1
- Phaeophyceae	2	-	-	1	-	1
- Ulvophyceae	1	-	-	-	-	1
Crustacea	8	-	2	3	-	3
- Amphipoda	2	-	-	1	-	1
- Cirripedia	1	-	-	1	-	-
- Copepoda	1	-	1	-	-	-
- Decapoda	3	-	-	1	-	2
- Diplostraca	1	-	1	-	-	-
Other taxa	19	1	8	1	4	5
- Actinopterygii	2	-	-	-	2	-
- Annelida	2	-	-	1	-	1
- Bryozoa	2	-	-	-	1	1
- Cnidaria	3	1	2	-	-	-

Taxon	<i>N</i>	NK	LO	PH	HI	SE
- Ctenophora	1	-	-	-	-	1
- Hexapoda	1	-	1	-	-	-
- Mollusca	4	-	3	-	-	1
- Nematoda	1	-	-	-	-	1
- Tunicata	3	-	2	-	1	-
Totals	37	1	12	7	4	13

*Table 2 . Pathways of introduction of marine alien species known to reproduce in Norway by 2018. Numbers are provided for all marine alien species, as well as split by ecological impact (NK–PH = "potentially high" or lower impact, HI+SE = "high" or "severe impact") and taxonomy. The number of the pathways in a row does not add up to the number of species (*N*) because a species may have more than one pathway. (Modified from Norwegian Biodiversity Information Centre , 2018).*

Subset	<i>N</i>	Release	Escape	Contaminant	Stowaway	Unaided
All introductions	37	2	4	4	27	19
By impact						
- NK–PH	20	-	1	2	16	11
- HI+SE	17	2	3	2	11	8
By taxonomy						
- Algae	10	-	1	-	10	5
- Crustaceans	8	1	1	2	6	5
- Other taxa	19	1	2	2	11	9

Table 3. Pathways of secondary spread of marine alien species known to reproduce in Norway by 2018. Information on secondary spread was unavailable or inapplicable for four species. See Table 2 for further explanations.

Subset	<i>N</i>	Release	Escape	Contaminant	Stowaway	Unaided
All sec. spread	33	1	-	2	12	24
By impact						
- NK–PH	17	-	-	1	5	12
- HI+SE	16	1	-	1	7	12
By taxonomy						
- Algae	7	-	-	-	3	7
- Crustaceans	8	-	-	1	3	7
- Other taxa	18	1	-	1	6	10

Table 4. Pathway subcategories of marine alien species known to reproduce in Norway by 2018. Pathways are listed for introduction

to, and secondary spread within, Norwegian nature; both for the entire set of marine alien species and for the subset with high or severe ecological impact ("HI+SE"). The number of the pathways in a column does not add up to the number of species because a species may have more than one pathway. (Modified from Norwegian Biodiversity Information Centre , 2018).

Pathway subcategory	Introduction		Secondary spread	
	All species	HI+SE	All species	HI+SE
Unaided – natural dispersal	21	8	26	13
Stowaway – ballast water	19	9	3	2
Stowaway – hull fouling	17	7	7	3
Stowaway – fishing equipment	9	5	1	1
Stowaway – others	2	1	2	2
Stowaway – hitchhikers on boats	1	1	1	1
Release – for fishery	2	2	1	1
Contaminant – parasites on animals	3	2	2	1
Contaminant – on animals (except par.)	2	1	-	-
Escape – from aquaculture	2	2	-	-
Escape – life food/bait	1	1	-	-
Escape – from aquaria	1	-	-	-
Totals (species)	37	17	33	16

Table 5. Pathways of introduction for the 13 marine alien species in Norway that have severe ecological impact. Pathway (sub)categories are unaided, stowaway (ballast water, hull fouling, fishing equipment, others, boats), release (for fishery), contaminant (parasites or other contaminants on animals) and escape (from aquaculture or of life food) (Modified from Norwegian Biodiversity Information Centre , 2018).

Species	Taxon. group	Un-aid.	Stowaway					Rel.	Cont.		Escape	
			BW	HF	FE	other	Boat	Fish	PA	CA	AC	LF
<i>Bonnemaisonia hamifera</i>	Algae	X	X	X	X							
<i>Codium fragile</i>	Algae	X	X	X								
<i>Dasysiphonia japonica</i>	Algae		X		X							
<i>Gracilaria vermiculophylla</i>	Algae		X		X		X					
<i>Sargassum muticum</i>	Algae	X			X	X						
<i>Caprella mutica</i>	Crustacea		X	X						X		
<i>Homarus americanus</i>	Crustacea											X
<i>Paralithodes camtschaticus</i>	Crustacea	X						X				
<i>Anguillicoloides crassus</i>	Nematoda								X			
<i>Crassostrea gigas</i>	Mollusca	X									X	
<i>Marenzelleria viridis</i>	Annelida	X	X									
<i>Mnemiopsis leidyi</i>	Ctenophora	X	X									
<i>Tricellaria inopinata</i>	Bryozoa			X								




Figure 49. Geographical distribution of all marine alien species in Norway. Numbers of alien reported from different Norwegian counties (plus Jan Mayen and Svalbard), for all species with an establishment category of C1 or higher (i.e. "surviving in the wild", $N = 39$; source: Sandvik et al., 2019).




Figure 50. Geographical distribution of invasive marine alien species in Norway. Numbers of alien reported from different Norwegian counties (plus Jan Mayen and Svalbard), only species with an establishment category of E (i.e. "dispersing, surviving and reproducing at multiple sites", N = 27; source: Sandvik et al., 2019).




Figure 51. Cumulative number of newly recorded marine alien species in Norway by year. Based on the 37 marine alien species known to reproduce in Norway by 2018 (Source: Artsdatabanken, 2018).

4.2 - Door knocker species to Norwegian waters

In addition to alien species reproducing in Norway, the Alien Species List contains 60 impact assessments of door knockers to Norwegian marine waters, 22 of which were categorised as having a high or severe ecological impact (Table 6). Information on pathways of introduction was available for 58 of the species. A total of 114 pathways of introduction were reported, which amounts to an average of 2.0 pathways per species (range 1–4). Again, stowaways and unaided spread were the dominant pathways of introduction, and pathways did not differ between taxonomic groups or impact categories (Table 7).

Subcategories of pathways are summarised in Table 8. As with alien species already present in Norway, the most important subcategories of stowaways were hull fouling, ballast water and fishing equipment.

Table 6. Ecological impact assessments of marine door-knockers to Norway risk-assessed in 2018. Tabulated by taxon and ecological impact category (NK = "no known impact", LO = "low impact", PH = "potentially high impact", HI = "high impact", SE = "severe impact"). (Source: Norwegian Biodiversity Information Centre , 2018).

Taxon	N	NK	LO	PH	HI	SE
Algae	12	3	2	5	-	2
Bonnemaisoniales	1	-	-	1	-	-
Ceramiales	4	1	1	2	-	-
Gigartinales	1	-	1	-	-	-
Halymeniales	2	-	-	2	-	-
Phaeophyceae	3	2	-	-	-	1
Ulvophyceae	1	-	-	-	-	1
Crustacea	20	-	8	3	6	3
Amphipoda	2	-	-	-	1	1
Cirripedia	3	-	2	-	1	-
Decapoda	8	-	3	1	2	2
Diplostraca	3	-	1	1	1	-
Poecilostomatoida	3	-	2	-	1	-
Pycnogonida	1	-	-	1	-	-
Other taxa	28	1	12	4	7	4
Actinopterygii	3	-	1	-	1	1
Annelida	3	-	1	1	1	-
Bryozoa	3	-	2	-	1	-
Cnidaria	3	1	2	-	-	-
Mollusca	11	-	5	2	2	2
Porifera	1	-	-	-	1	-
Trematoda	1	-	1	-	-	-
Tunicata	3	-	-	1	1	1
Totals	60	4	22	12	13	9

Table 7. Pathways of introduction of marine door-knocker species to Norway risk-assessed in 2018. None of the species was expected to be introduced by means of release or escape. Numbers are provided for all species, as well as split by ecological impact (NK–PH = "potentially high" or lower impact, HI+SE = "high" or "severe impact") and taxonomy. The number of the pathways in a row does not add up to the number of species (N) because a species may have more than one pathway. (Modified from Norwegian Biodiversity Information Centre , 2018).

Subset	N	Contaminant	Stowaway	Unaided
All introductions	58	10	48	30
By impact				
NK–PH	37	8	30	21
HI+SE	21	2	18	9
By taxonomy				
Algae	12	3	12	7
Crustaceans	20	3	16	12
Mollusca	9	2	6	4
Other taxa	17	2	14	7

Table 8. Pathway subcategories of marine door-knocker species to Norway risk-assessed in 2018. Pathways are listed both for the entire set of door-knockers and for the subset with high or severe ecological impact ("HI+SE"). The number of the pathways in a column does not add up to the number of species because a species may have more than one pathway. (Modified from Norwegian Biodiversity Information Centre , 2018).

Pathway subcategory	Introduction	
	All species	HI+SE
Unaided – natural dispersal	29	9
Stowaway – ballast water	20	10
Stowaway – hull fouling	41	14
Stowaway – fishing equipment	14	5
Contaminant – parasites on animals	5	1
Contaminant – on animals (except par.)	5	1
Totals (species)	58	21

For fish, so far there are only two established species relevant for marine ecosystems, however there are several door knockers and potential door knockers which deserves attention (Table 9). The Norwegian Biodiversity Information Centre has in 2021 identified potential door knockers, which are going through a horizon scanning for possible inclusion in the new risk assessment starting in 2022. The door knocker associated with the highest risk so far is the round goby, *Neogobius melanostomus* (Artsdatabanken 2018). The most likely dispersal pathways for this species are unaided dispersal from Sweden and Denmark and ship traffic (ballast water and possibly eggs deposited on ships' hulls). Notably, many of the potential door knockers are gobies from the Ponto-Caspian region (Black Sea, Caspian Sea, Azov Sea). These are spread via ballast water,

and in continental Eurasia also by corridors. After being introduced to new areas, they often have a high potential for unaided secondary dispersal.

For Svalbard, the door knocker pink salmon, *Oncorhynchus gorbuscha* (Table 10), came unaided from Russia and is already present in the sea around Svalbard but is not reproducing. Increasing temperatures may make reproduction possible in the future (in fresh water).

Shipping, together with unaided dispersal, are the most likely vectors for new introductions of alien marine fish to Norway. Ballast water and hulls in ships can transport fish of all life stages. An adult blenny, *Parablennius zvonimiri*, was for example found in western Norway inside a barnacle shell on a ship from the Mediterranean a few years ago. Occasional findings of new fish species introduced by ships do occur, but so far these do not seem to have established (Table 9). A review showed that fish found in ballast tanks most often were from the families Gobiidae and Clupeidae. On lists of ballast mediated introductions, Gobiidae, Blenniidae and Pleuronectidae was frequent, and gobies and blennies were the families that most often managed to establish (Wonham et al. 2000).

Table 9. Marine, anadromous, catadromous and amphidromous alien fish (Actinopterygii) of relevance for Norway, their associated risk, occurrence and main vectors.

Species	Risk ¹	Native Area	Occurrence in Norway or closest occurrence as NIS	Main vector(s) for introduction to Norway
¹Established alien species:				
Pink salmon, <i>Oncorhynchus gorbuscha</i>	HI	N Pacific, Arctic	yes	Unaided
Rainbow trout, <i>Oncorhynchus mykiss</i>	HI	N Pacific, NW America	yes	Escape, release, unaided
¹Door-knockers:				
American eel, <i>Anguilla rostrata</i>	HI	W Atlantic	Germany(?)	Unaided
Round goby, <i>Neogobius melanostomus</i>	SE	Ponto-Caspian	Sweden	Ship traffic, unaided
False kelpfish, <i>Sebastes marmoratus</i>	NR	Asia, Oceania	Fredrikstad 2016	Ship traffic (ballast water)
Atlantic croaker, <i>Micropogonias undulatus</i>	LO	W Atlantic	Denmark	Ship traffic (ballast water), unaided
²Potential door knockers (selection):				
<i>Babka gymnotrachelus</i> , racer goby	-	Ponto-Caspian	Germany, Poland	Ship traffic (ballast water)
<i>Coregonus peled</i> , peled	-	e.g. Russia	Germany, Poland	Release
<i>Fundulus heteroclitus</i> , mummichog	-	W Atlantic, NE America	Portugal, Spain	Release, escape
<i>Gobiosoma bosc</i> , naked goby	-	W Atlantic	Netherlands, Germany	Ship traffic, unaided
<i>Morone americana</i> , white perch	-	W Atlantic, NE America	-	Aquarium? Release?
<i>Neogobius fluviatilis</i> , monkey goby	-	Ponto-Caspian	Netherlands, Poland	Ship traffic, unaided
<i>Oncorhynchus kisutch</i> , coho salmon	-	N Pacific	Belgium	Unaided
<i>Ponticola kessleri</i> , big head goby	-	Ponto-Caspian	Netherlands, Germany	Ship traffic, unaided
<i>Proterorhinus marmoratus</i> , tubenose goby	-	Ponto-Caspian	Netherlands, Germany	Ship traffic, unaided
<i>Sphaeroides pachygaster</i> , blunthead puffer	-	Circumglobal, nearest: UK	Mediterranean	Ship traffic (ballast water)
<i>Vimba vimba</i> , vimba bream	-	Eurasia	Netherland	Release
³Occasional finding of potential door knockers (selection):				

Species	Risk	Native Area	Occurrence in Norway or closest occurrence as NIS	Main vector(s) for introduction to Norway
<i>Glossanodon leioglossus</i> , small toothed argentine	NR	E Atlantic, Mediterranean	Sognefjord 1942	Ship traffic (ballast water)
<i>Parablennius zvonimiri</i>	NR	Mediterranean, Black Sea	Karmøy 2017	Ship traffic
<i>Scartella cristata</i> , Molly miller	-	W & E Atlantic, Mediterranean	Karmøy 2016	Ship traffic

¹From the Alien Species List of Norway (Artsdatabanken, 2018). Ecological risk assessment, SE=severe impact, HI= high impact, LO=low impact, NR=not risk assessed.

²Selected fish species of interest from the ongoing horizon scanning by the Norwegian Biodiversity Information Centre (unpubl., autumn 2021).

³Occasional findings of nonindigenous marine fish in Norway, from the Alien Species List of Norway (2018) and Norwegian Institute of Marine Research

Table 10. Marine and anadromous alien fish (*Actinopterygii*) relevant for Svalbard, geographical occurrence and likely vectors for introduction.

Species	Native Area	Occurrence Europe/Norway/Svalbard	Main vector(s)
¹Door knocker:			
Pink salmon, <i>Oncorhynchus gorbuscha</i>	N Pacific, Arctic	yes / yes / yes	unaided
²Potential door knockers:			
Yellowfin goby, <i>Acanthogobius flavimanus</i>	NW Pacific, Asia	no / no / no	shipping, ballast water
Giant grenadier, <i>Albatrossia pectoralis</i>	N Pacific	no / no / no	ballast water
European flounder, <i>Platichthys flesus</i>	N Atlantic	yes / yes / no	ballast water
Coho salmon, <i>Oncorhynchus kisutch</i>	N Pacific	yes / no / no	unaided, release?
Rainbow trout, <i>Oncorhynchus mykiss</i>	NW America	yes / yes / no	unaided, release?

¹ From the Alien Species List of Norway (Norwegian Biodiversity Information Centre, 2018). Likely to establish with warmer temperatures.

² Mentioned in online resource to van den Heuvel-Greve et al. (2021)

5 - Management of dispersal vectors and high impact species

5.1 - Measures for prevention of introduction and dispersal of NIMS

Preventive measures early in the process of biological invasions are more cost-efficient than control measures later (Leung et al. 2002). If prevention fails, early detection and rapid response are important in fighting alien species (Hulme 2006, Lehtiniemi et al. 2015). There are many different types of action to take from securing international collaboration and agreements, national monitoring and surveys at high-risk sites, to local eradication measures (e.g. Lehtiniemi et al. 2015, Table 11). Also, one should not forget the importance of maintaining healthy well-functioning ecosystems not too much affected by human impacts, which likely are more robust to invasion of alien species. Predators have an important role and biological control by natural predators should help keep the density of alien species down (Madenjian et al. 2011, Ojaveer et al. 2015).

Table 11. Process for biological invasion, main dispersal pathways, and possible measures to reduce the effect of alien marine species.

Process for biological invasion	Dispersal pathway	Goal of measure	Example of measure
Transport, arrival, introduction	Shipping <ul style="list-style-type: none"> • Ballast water • Fouling • Escape cultivation • Aquaculture Release • Live sea food • Aquarium Unaided 	Prevent introduction, precautionary approach	-Legislation and international agreements (ballast water and sediments, biofouling, trade in live organisms) -closed cages and/or aquaculture on land - public awareness
Establishment		Early detection, rapid response, eradication	-Predicting hotspots for introduction -Port monitoring -Surveys of offshore infrastructure -Inspecting boat hulls -Public awareness and citizen science - Catching -Healthy ecosystem
Secondary spread	Unaided Ship and boat traffic	Limit and control	-Marina surveys -Catching/removal -Removing hull fouling -Ballast water management
Impact on ecosystems		Mitigation of impact	-Catching -Removal in especially vulnerable areas

In Norway several measures for preventing NIMS entering coastal ecosystems is already in place such as ballast water treatment and regulations on import of organisms for aquaculture or food.

Even though it is stated by law that it is the sellers of alien species for food or aquarium duty to inform about the risk associated with specimens or larvae accidentally released to the environment, there is a certain uncertainty whether this system is functioning well enough. Increased information to traders on this duty is recommended.

Biofouling on vessels is then probably the vectors with the highest risk for introduction of NIMS into the Norwegian coast currently. There is currently ongoing work in IMO to evaluate experience from the guideline and develop standards for cleaning and control of vessels. As a part of this work an industrial standard for in-water cleaning with capture is under development (Anon 2021). This work will most likely take years before

measures are implemented internationally. In the mean while it is recommended to give more information and encourage the ships owners to keep their vessels as clean as possible. Regional regulations on vessel traffic out from port with *Didemnum vexillum* should be considered.

Long distance transport of living cleaner fish and release of transport water into the environment is not recommended (Mortensen et al. 2021).

The increasing problem with floating litter and hitch hikers on this cannot be solved without a worldwide change in human behaviour to decrease this flow.

5.2 - Measures for eradication of problematic NIMS

Due to the lack of systematic monitoring and mapping programmes of NIMS in Norway is it often too late for eradication when a problematic species first is discovered. To be able to make a quick and determined eradication attempt one relies on an early warning system based on regularly monitoring of coastal sites. Moreover, there is a need for an alert group consisting of relevant management directories and researchers to decide on which actions it is appropriate to take. Such a group could have a sleeping funding system, which quickly allows actions in form of mapping of the abundance and dispersal of the species and eradication attempts. The need for such an action group is highlighted by the example of the newly introduced species *Didemnum vexillum* in Norway where it has taken time to establish contact between relevant authorities and to date no sizable measures have been taken to minimize the spread between ports one year after the first record.

5.2.1 - Pathogens

Pathogens may be spread via translocation of farmed species, release of non-indigenous species or live seafood meant for human consumption. Once established in a marine species, pathogens are difficult or impossible to eradicate. Pathogens of farmed animals are listed according to their severity / effect on the susceptible host species. Detection of such notifiable diseases require a response, and mitigation measures are establishment of a protection/eradication zone and removal of infected stock. For some notifiable diseases there are national surveillance programmes. Other diseases may be detected during health inspections or control. For wild animals there is no such surveillance. To minimize risk of disease introductions and transmission, long-distance translocations of live animals with un-known health status should, as a general rule, be avoided.

5.2.2 - Pacific oysters

After the bio-invasion of Pacific oysters, commencing in 2007 (Wrange et al. 2010), mitigation was discussed, and measures were taken to remove oysters from beaches and areas used by the public. Pacific oysters settle on shallow water and may build dense populations. The oysters have sharp edges and stepping on them may cause severe wounds. Removal is organized every summer, as local campaigns. The campaigns however remove only a small fraction of the total amount of oysters. They help to protect users of beaches and other public areas but have no effect on the further spread of oysters along the coast. Pacific oysters must be regarded as a new species, permanently established in the Norwegian fauna. We consider it important to discuss the mitigation and management issues and turn the situation from being a problem into also being a resource (Mortensen et al 2019). We have therefore assessed the risk and studied the effect on affected habitats, thus providing background for mitigation measures as well as management of the resource (Mortensen et al. 2017; 2019). The 2019-report proposes a process focusing on a new management model, combining exploitation with removal. We are now carrying out studies on the establishment in different habitats in the invasion front (south and north).

5.2.3 - Pink salmon

There has been a dramatic invasion of pink salmon in Norway every second year from 2017, especially in the north. The number of registered pink salmon increased from 2017 to 2019 (Berntsen et al. 2018, 2020). There has been targeted removal of pink salmon in designated rivers by intense angling and/or fishing with nets or traps, from June to September all three years (2017, 2020, 2021). In addition, sports fishing in rivers and bag- or bend-net fishing in the sea has caught a substantial number of pink salmon. The activity was accomplished thanks to the Norwegian Environmental Agency, the County governor, several Norwegian institutions and research institutes, and voluntary efforts. Removal of pink salmon in rivers summertime is a good and necessary measure to mitigate the effects of the species as much as possible. The next expected invasion is in 2023, and increased resources are needed to remove as many pink salmon as possible.

5.2.4 - *Didemnum vexillum*

Several studies have been performed to find the best method to destroy *D. vexillum* when growing on mussel and oyster cultures without impacting the mollusc. Different chemical solutions such as acetic acid, lime, chlorine, sodium hydroxide and bleach as well as brine, fresh water and desiccation. Lime solutions (3-4%), sodium hydroxide (0,5) and acetic acid (5%) effectively killed *D. vexillum* in less than 10 minutes, but acetic acid and sodium hydroxide had a severe impact on the molluscs. The use of brine and freshwater needed four hours to kill *D. vexillum* (McKenzie et al. 2017 and references therein). Testing out different chemical solutions in special made enclosed berths for removal of biofouling species on recreational vessels showed that acetic acid and sodium hydroxide efficiently removed the biota from the vessels (Roche et al. 2015).

Due to the combined sexual and clonal reproduction of the species, it has proved extremely hard to eradicate *D. vexillum* from natural sites even if the distribution of colonies has been limited. A massive eradication attempt was tried out in New Zealand in the period 2003-2009 by cleaning of all vessels, capsuling pilings and artificial structures with plastic sheeting and covering up the seafloor with new sediment. The operation was costly and removed the species for a short time, but new biomass of the species built up rapidly the following year. A similar attempt was made in Holyhead Marina, UK with the same results (McKenzie et al. 2017 and references therein). *D. vexillum* is now spread to five ports in Norway and new colonies are reported frequently in the vicinity of the ports, and a costly eradication attempt is likely to fail.

As ascidians have a limited natural dispersal capacity due to the larval settlement within 24 hours after spawning, measures to prevent/slow down further human aided dispersal can be taken. Such measures can be for example cleaning of vessels that has been laying for a period (>24 hours in summer (April-October), 14-30 days in winter) in ports which is infected before leaving the port. Cleaning of fishing gear, instruments etc which have been in the sea for more than 24 hours is recommended, by drying or freshwater treatment in more than 4 hours. *D. vexillum* is yet to be detected on Norwegian salmon farms but will probably cause large problems if fouling the nets. The usual praxis of flushing the nets frequently during the summer season will likely cause further spread of the species, as small fragments of the colonies grow rapidly into new colonies. Likewise, will the species potential cause problems for mussel farms and kelp farms by overgrowing and smothering of the aquaculture organisms. More research is needed to find the best protocol for early detections and removal of *D. vexillum* at aquaculture facilities.

5.2.5 - Round goby

The round goby, a door knocker with severe impact, can serve as an example of measures to handle potential introductions of new alien marine fish species dispersing unaided or by shipping. The round goby is a brackish species with a broad salinity tolerance (Kornis et al. 2012). In a spatial analysis based on the main dispersal pathways together with environmental preferences of the species (Kotta et al. 2016), hotspots for likely introduction and establishment were modelled (Forsgren & Hanssen, unpubl.). Shallow, brackish areas near

large international ports, especially in the Oslofjord region, came out as the most likely areas for introduction and establishment. Since early detection is crucial for efficient mitigation, yearly monitoring of selected ports by means of eDNA sampling would be a good strategy. Efforts to try and catch fish with passive or active fishing gear may be useful in addition. After establishment, the species have a high potential to disperse and further expanding its range. Range expansion was in Denmark estimated to about 30 km per year (Azour et al. 2015). In the Baltic, eradication of the round goby is unrealistic but several measures to limit impact and spread has been suggested, including monitoring and management of ships' ballast water and hull fouling (Ojaveer et al. 2015). The round goby can also disperse from the sea into freshwater, as found in the Baltic where it now has invaded several salmonid streams (Verliin et al. 2017). In North America, electrical barriers in rivers to stop dispersal have been tried and proved efficient, in addition to baited traps and other measures (reviewed in Kornis et al 2012, Forsgren & Florin 2018).

6 - References

- Agnalt A-L, Jørstad KE, Pavlov, Farestveit E, Sundet J. 2011. The snow crab, *Chionoecetes opilio* (Decapoda, Majoidae, Oregoniidae) in the Barents Sea. In: In the Wrong Place: Alien Marine Crustaceans—Distribution, Biology and Impacts. Invading Nature, Springer Series in Invasion Ecology, 6, pp. 283–300. Ed. by B. S. Galil, P. F. Clark, and J. T. Springer Science and Business Media Carlton, London.
- Ahne W. 1980. Experimental infection of pike (*Esox lucius*) with Egtvedvirus. Tierarztl Umsch 35: 225-229.
- Alvsvåg J, Agnalt A-L, Jørstad KE. 2009. Evidence for a permanent establishment of the snow crab (*Chionoecetes opilio*) in the Barents Sea. Biological Invasions, 11, 587-595.
- Anon. 2015. Forskrift om fremmede organismer. LOVDATA. FOR-2015-06-19-716.
- Anon. 2021. Industry standard on in-water cleaning with capture. BIMCO. Ver.1.0. <https://www.bimco.org/about-us-and-our-members/publications/industry-standard-on-in-water-cleaning-with-capture>
- Artsdatabanken. 2018. Fremmedartslista 2018, Trondheim. <https://www.artsdatabanken.no/fremmedartslista2018>
- Azour F, van Deurs M, Behrens J, Carl H, Hüsey K, Greisen K, Ebert R, Møller PR. 2015. Invasion rate and population characteristics of the round goby *Neogobius melanostomus*: effects of density and invasion history. Aquatic Biology 24:41-52.
- Barrett, CJ, A Cook, D Stone, C Evans, D Murphy, P Johnson, M Thain, G Wyn, M Grey, H Edwards, D Quigley, Stebbing PD. 2020. A review of American lobster (*Homarus americanus*) records around the British Isles: 2012-2018. Hydrobiologia, 847: 3247-3255.
- Berntsen HH, Sandlund OT, Ugedal, O, Thorstad E, Fiske P, et al. 2018. Pukkellaks i Norge, 2017. NINA Rapport 1571. Norsk institutt for naturforskning. <http://hdl.handle.net/11250/2575646>
- Berntsen HH, Sandlund OT, Thorstad EB, Fiske P. 2020. Pukkellaks i Norge, 2019. NINA Rapport 1821. Norsk institutt for naturforskning. <https://hdl.handle.net/11250/2651741>
- Blackburn TM, Pyšek P, Bacher S, Carlton JT, Duncan RP, Jarošík V, Wilson JR, Richardson DM. 2011. A proposed unified framework for biological invasions. Trends in Ecology and Evolution 26: 333–339. <http://dx.doi.org/10.1016/j.tree.2011.03.023>
- Blanco Gonzalez E, Knutsen H, Jorde PE. 2016. Habitat discontinuities separate genetically divergent populations of a rocky shore marine fish. PLoS ONE 11. <https://doi.org/10.1371/journal.pone.0163052>
- Boersma M, Malzahn A M, Greve W, et al. 2007. The first occurrence of the ctenophore *Mnemiopsis leidyi* in the North Sea, Helgol. Mar. Res., 61:153-155
- Brockhoff A, McLay C. 2011. Human-mediated spread of alien crabs. In: Galil BS, Clark PF, Carlton JT (eds) In the wrong place—Alien marine crustaceans: distribution, biology and impacts, vol 6. Invading Nature—Springer Series in Invasion Ecology. pp 27–106. https://doi.org/10.1007/978-94-007-0591-3_2
- Campell ML, King S, Heppenstall LD, van Gool E, Martin R, Hewitt CH. 2017. Aquaculture and urban marine structures facilitate native and non-indigenous species transfer through generation and accumulation of marine debris. Marine Pollution Bulletin 123: 304-312.

- Carlton, JT. 1985. Transoceanic and interoceanic dispersal of coastal marine organisms: the biology of ballast water. *Oceanogr. Mar. Biol. Ann. Rev.* 23: 313-371.
- Carlton JT, Chapman JW, Geller JB, Miller JA, Carlton DA, McCuller MI, Treneman NC, Steeves BP, Ruiz G. 2017. Tsunami-driven rafting: Transoceanic species dispersal and implications for marine biogeography. *Science* 357: 1402-1406.
- CBD. 2014. Pathways of introduction of invasive species, their prioritization and management. Convention on Biological Diversity, Subsidiary Body on Scientific, Technical and Technological Advice, Montréal.
<https://www.cbd.int/doc/meetings/sbstta/sbstta-18/official/sbstta-18-09-add1-en.pdf>
- Costello JH, Bayha KM, Mianzan HW, Shiganova TA, Purcell JE. 2012. Transitions of *Mnemiopsis leidyi* (Ctenophora: Lobata) from a native to an exotic species: a review. *Hydrobiologia* 690: 21-46
- Dahle G, Sainte-Marie B, Mincks SL, Farestveit E, Jørstad KE, Hjelset AM, Agnalt A-L. Submitted. Genetic analysis of the exploited snow crab (*Chionoecetes opilio*) in the Barents Sea – revealing pathway of introductions. *ICES J. Mar. Sci.*
- Danielsen HEH, Hjelset AM, Bluhm BA, Hvingel C, Agnalt A-L. 2019. A first fecundity study of the female snow crab *Chionoecetes opilio* Fabricius, 1788 (Decapoda: Brachyura: Oregoniidae) of the newly established population in the Barents Sea. *J Crust Biol.* <https://doi.org/10.1093/jcbiol/ruz039>[Furlani 1996.](https://doi.org/10.1093/jcbiol/ruz039)
- Darling JA, Bagley MJ, Roman J, Tepolt CK, Gellers JB. 2008. Genetic patterns across multiple introductions of the globally invasive crab genus *Carcinus*. *Molecular Ecology*, 17: 4992-5007.
- Dionne M, Sainte-Marie B, Bourget D, Gilbert D. 2003. Distribution and habitat selection of early benthic stages of snow crab *Chionoecetes opilio*. *Mari. Ecol. Prog. Ser.*, 259: 117-128.
- Dybedal P. 2018. Cruise passenger traffic to Norway-history and forecasts 2018-2060. Report nr 1651/2018. Norwegian centre for transport research.
- Erkinharju T, Dalmo RA, Hansen M, Seternes T. 2020. Cleaner fish in aquaculture: review on diseases and vaccination. *Reviews in Aquaculture*. 1–49. <https://doi.org/10.1111/raq.12470><https://doi.org/10.1111/raq.12470>
- Faust E, André C, Meurling S, Kochmann J, Chritiansen H, Jensen LF, Charrier G, Laugen AT, Strand Å. 2017. Origin and route of establishment of the invasive Pacific oyster *Crassostrea gigas* in Scandinavia. *Marine Ecology Progress Series* 575:95-105. <http://www.int-res.com/abstracts/meps/v575/p95-105/>
- Faust E, Halvorsen KT, Andersen P, Knutsen H, André C. 2018. Cleaner fish escape salmon farms and hybridize with local wrasse populations. *Royal Society Open Science* 5: 171752.
<https://doi.org/10.1098/rsos.171752>
- Forsgren E, Florin A-B. 2018. Svartmunnet kutling – en invaderende fremmed fiskeart på vei mot Norge. NINA Rapport 1454. Norsk institutt for naturforskning. <http://hdl.handle.net/11250/2479873>
- Fossøy F, Sivertsgård R, Ambjørndalen VM, Brandsegg H, Andersskog IPØ, Husa V, Forsgren E. 2022. Kartlegging av den fremmede marine arten havnespy *Didemnum vexillum* ved hjelp av miljø-DNA. En rask respons undersøkelse. NINA Rapport 2092. Norsk institutt for naturforskning.
- Goldsmith J, Nudds SH, Stewart DB, Higdon JW, Hannah CG, Howland KL. 2018. Where else? Assessing zones of alternate ballast water exchange in the Canadian eastern Arctic. *Mar Pollut Bull.* 139:74–90.

- Gollasch S. et al. 2002. Life in Ballast Tanks. In: Leppäkoski E., Gollasch S., Olenin S. (eds) Invasive Aquatic Species of Europe. Distribution, Impacts and Management. Springer, Dordrecht. https://doi.org/10.1007/978-94-015-9956-6_23
- Govindarajan AF, Källström B, Selander E, Östman C, Dahlgren TG. 2019. The highly toxic and cryptogenic clinging jellyfish *Gonionemus* sp. (Hydrozoa, Limnomedusae) on the Swedish west coast. PeerJ. 7:e6883. doi: 10.7717/peerj.6883
- Granhag L, Møller LF, and Hansson LJ 2011. Size-specific clearance rates of the ctenophore *Mnemiopsis leidyi* based on in situ gut content analyses. Journal of Plankton Research 33,: 1043–1052.
- Grefsrud ES, Karlsen Ø, Kvamme BO, Glover K, Husa V, Hansen PK, Samuelsen OB, Sandlund N, Stien LH, Svåsand T. 2021a. Risikoreport norsk fiskeoppdrett - risikovurdering (in Norwegian), Rapport fra Havforskningen, 2021-8, 198 pp. <https://www.hi.no/hi/nettrapporter/rapport-fra-havforskningen-2021-8>
- Grefsrud ES, Karlsen Ø, Kvamme BO, Glover K, Husa V, Hansen PK, Samuelsen OB, Sandlund N, Stien LH, Svåsand T. 2021b. Risikoreport norsk fiskeoppdrett 2021 - kunnskapsstatus (in Norwegian), Rapport fra Havforskningen, 2021-7, 280 pp.
- Gregory M. 2009. Environmental implications of plastic debris in marine settings-entanglement, ingestion, smothering, hangers-on, hitch-hiking and alien invasions. Philosophical Transactions 364:2013-2025.
- Grotmol S, Totland GK, Thorud K, Hjeltnes BK. 1997. Vacuolating encephalopathy and retinopathy associated with a nodavirus-like agent: a probable cause of mass mortality of cultured larval and juvenile Atlantic halibut *Hippoglossus hippoglossus*. Diseases of Aquatic Organisms 29: 85–97.
- Guðmundsdóttir S, Vendramin N, Cuenca A, Sigurðardóttir H, Kristmundsson A, Moesgaard Iburg T, Jørgen Olesen N 2019. Outbreak of viral haemorrhagic septicaemia (VHS) in lumpfish (*Cyclopterus lumpus*) in Iceland caused by VHS virus genotype IV. Journal of Fish Diseases 42: 47–62. <https://doi.org/10.1111/jfd.12910>
- Gulliksen B. 1971. A new record of *Gonionemus vertens* Agassiz (Limnomedusea) in Norway. K. norske Vidensk. Selsk. Skr. 1971. 1-4
- Gündoğdu S, Çevik C, Karaca S. 2017. Fouling assemblage of benthic debris collected from Mersin Bay, NE Levantine coast of Turkey. Marine Pollution Bulletin 124: 147-154.
- Hallegraeff GM. 1993. A review of harmful algal blooms and their apparent global increase* Phycologia (1993) Volume 32 (2)
- Hansen HSB. 2016. Three major challenges in managing non-native sedentary Barents Sea snow crab (*Chionoecetes opilio*). Marine Policy, 71: 38-43.
- Hansen C, Mousing EA, Hvingel C. In prep. Snow crab in the Barents Sea, winner or villain?
- Hansson HG. 2006. Ctenophores of the Baltic and adjacent seas - the invader *Mnemiopsis* is here! Aquatic Invasions 1, Issue 4: 295 -298
- Haraldstad T, Hindar A, Hobæk A, Håvardstun J, Skjelbred B. 2013. Resipientundersøkelse av Reddalsvann, Grimstad. NIVA Rapport L.NR. 6485-2013.
- Harrower CA, Scalera R, Pagad S, Schönrogge K, Roy HE. 2020. Guidance for interpretation of the CBD categories of pathways for the introduction of invasive alien species. Publications Office of the European Union,

Luxembourg. <http://dx.doi.org/10.2779/6172>

Hess-Erga OK, Moreno-Andrés J, Enger Ø, Vadstein O. 2019. Microorganisms in ballast water: disinfection, community dynamics, and implications for management. *Science of the Total Environment* 657, 704-716

Heuvel-Greve MJ, van den Brink AM, Glorius ST, de Groot A, Laros I, Renaud PE, Pettersen R, Weslawski JM, Kutlinski P, Murk AJ. 2021. Early detection of marine non-indigenous species on Svalbard by DNA metabarcoding of sediment. *Polar Biology* 44:653-665.

Hjelset AM. 2012. Female life-history parameters in the introduced red king crab (*Paralithodes camtschaticus*, Tilesius 1815) in the Barents Sea: A study of temporal and spatial variation in three Norwegian fjords. PhD. Faculty of Biosciences, Fisheries and Economics. Department of Arctic and Marine Biology, University of Tromsø, Norway.

Hjelset AM, Danielsen HEH, Westgaard JI, Agnalt A-L. 2021. Taxonomic and genetic confirmed findings of snow crab (*Chionoecetes opilio*) larvae in the Barents Sea. *Polar Biology* 44:2107-2115.

Holt RE, Hvingel C, Agnalt A-L, Dolgov AV, Hjelset AM, Bogstad B. 2021. Snow crab (*Chionoecetes opilio*), a new food item for North-east Arctic cod (*Gadus morhua*) in the Barents Sea. *ICES J. Mar. Sci.* 78: 491-501.

Holte B, Fuhrmann MM, Tandberg AHS, Hjelset (AM). In prep. New data on infaunal and epifaunal benthic secondary production in the Barents Sea, with focus on snow crab (*Chionoecetes opilio*) consumption

Hosia A, Falkenhaug T. 2015. Invasive ctenophore *Mnemiopsis leidyi* in Norway. *Marine Biodiversity Records* 8: 1-9

Hulme PE. 2006. Beyond control: wider implications for the management of biological invasions. -*Journal of Applied Ecology* 43:835–847.

Hulme PE, Bacher S, Kenis M, Klotz S, Kühn I, Minchin D. et al. 2008. Grasping at the routes of biological invasions: a framework for integrating pathways into policy. *Journal of Applied Ecology* 45: 403–414.
<http://dx.doi.org/10.1111/j.1365-2664.2007.01442.x>

Husa V, Sjøtun KI. 2006. Vegetative reproduction in «*Heterosiphonia japonica*» (Dasyaceae, Ceramiales, Rhodophyta) an introduced red alga on European coasts. *Botanica Marina* 49: 191-199.

Husa V, Heggøy E, Sjøtun K, Agnalt AL, Johansen PO, Glenner H, Hatlen K. 2012 a. Kartlegging av fremmede marine arter i Hordaland. Utredning for DN 2-2012. Direktoratet for naturforvaltning.

Husa V, Heggøy E, Agnalt A-L, Sjøtun K, Svensen R, Rokkan-Iversen K, Alvestad T. 2012 b. Kartlegging av fremmede marine arter i Rogaland. Utredning for DN 3-2012. Direktoratet for naturforvaltning.

Husa V, Agnalt AL, Svensen R, Rokkan Iversen K, Steen H, Jelmert A, Farestvedt E, Petersen H. 2013. Kartlegging av fremmede marine arter i indre og ytre Oslofjord. Utredning for DN 4-2013. Direktoratet for naturforvaltning

Husa V, Agnalt AL, Svensen R, Floerl O, Brattegard T, Rokkan-Iversen K, Glenner H, Bruntveit L. 2014. På jakt etter fremmede marine arter i Narvik havn. Rapport fra Miljødirektoratet M222-2014

Huserbråten, M, Hvingel C, Danielsen HEH, Agnalt AL, Hjelset AM. In prep. Understanding the early life history of the invading snow crab (*Chionoecetes opilio*) in the Barents.

- Hwang J, Park SY, Lee S, Lee T-K. 2018. High diversity and potential translocation of DNA viruses in ballast water, *Marine Pollution Bulletin*, Volume 137, 2018, P .449-455,
ISSN 0025-326X, <https://doi.org/10.1016/j.marpolbul.2018.10.053.0>
- Jansson E, Quintela M, Dahle G, Albretsen J, Knutsen H, André C, Strand Å, Mortensen S, Taggart JB, Karlsbakk E, Kvamme BO, Glover KA. 2017. Genetic analysis of goldsinny wrasse reveals evolutionary insights into population connectivity and potential evidence of inadvertent translocation via aquaculture. *ICES Journal of Marine Science* 74: 2135–2147. <https://doi.org/10.1093/icesjms/fsx046>
- Jansson E, Besnier F, Malde K, André C, Dahle G, Glover KA. 2020. Genome wide analysis reveals genetic divergence between Goldsinny wrasse populations. *BMC Genetics* 21(1): 118. <https://doi.org/10.1186/s12863-020-00921-8>
- Jaspers C, Huwer B, Antajan E, et al. 2018. Ocean current connectivity propelling the secondary spread of a marine invasive comb jelly across western Eurasia. *Global Ecol Biogeogr.* 27: 814– 827. <https://doi.org/10.1111/geb.12742>
- Javidpour J, Sommer U, Shiganova T A, et al. 2006. First record of *Mnemiopsis leidyi* A. Agassiz 1865 in the Baltic Sea, *Aquatic Invasions*. vol. 1 (pg. 299-302)
- Jørstad, KE, E Farestveit, H Rudra, A-L Agnalt, Olsen S. 2002. Studies on red king crab (*Paralithodes camtschaticus*) introduced to the Barents Sea, pp. 425-438. In: *Crabs in Cold Water Regions: Biology, Management, and Economics*, AJ Paul, EG Dawe, R Elnor, GS Jamieson, GH Kruse, RS Otto, B Sainte-Marie, TC Shirley 6 D Woodby (eds). 19th Lowell Wakefield Fisheries Symposium. Alaska Sea Grant College Program. AK-SG-02-01. 866 p.
- Jørstad K, Agnalt AL, Farestveit E. 2011. The Introduced American Lobster, *Homarus americanus* in Scandinavian Waters. In (Vol. 6, pp. 625-638).
- Kaiser BA, Kourantidou M, Fernandez L. 2018. A case for the commons: The snow crab in the Barents: *Journal of Environmental Management*, 210: 338-348.
- Kiessling T, Gutow L, Thiel M. 2015. Marine Litter as Habitat and Dispersal Vector. In: Bergmann M., Gutow L., Klages M. (eds) *Marine Anthropogenic Litter*. Springer, Cham.
- Kinlan BP, Gaines SD. “Propagule Dispersal in Marine and Terrestrial Environments: A Community Perspective.” *Ecology*, vol. 84, no. 8, Ecological Society of America, 2003, pp. 2007–20, <http://www.jstor.org/stable/3450025>
- Knutsen H, Jorde PE, Gonzalez EB, Robalo J, Albretsen J, Almada V. 2013. Climate Change and Genetic Structure of Leading Edge and Rear End Populations in a Northwards Shifting Marine Fish Species, the Corkwing Wrasse (*Symphodus melops*). *PLoS ONE* 8, e67492. <https://doi.org/10.1371/journal.pone.0067492>
- Kornis MS, Mercado-Silva N, Vander Zanden MJ. 2012. Twenty years of invasion: a review of round goby *Neogobius melanostomus* biology, spread and ecological implications. *Journal of Fish Biology* 80:235-285.
- Korsnes K, Karlsbakk E, Skaar CK, Sælemyr L, Nylund A, Kvamme BO, Mortensen S. 2017. High nervous necrosis virus (NNV) diversity in wild wrasse (Labridae) in Norway. *Diseases of Aquatic Organisms* 126, 43–50. <https://doi.org/10.3354/dao03159>
- Kotta J, Nurkse K, Puntila R, Ojaveer H. 2016. Shipping and natural environmental conditions determine the

- distribution of the invasive non-indigenous round goby *Neogobius melanostomus* in a regional sea. Estuarine, Coastal Shelf Science 169:15-24.
- Kramp PL. 1922. "Demonstration of *Gonionemus murbachii* from Kristianiafjord, Norway." Vidensk. Medd. dansk. naturh. Foren. Kbh. (Meeting-report) 74: xi
- Kuzmin S, Olsen S. 1994. Barents Sea king crab (*Paralithodes camtschatica*). The transplantation experiments were successful. ICES C.M. 1994/K:12.
- Kuzmin S, Olsen S, Gerasimova O. 1996. Barents Sea king crab (*Paralithodes camtschatica*): The transplantation experiments were successful, pp. 649-663. In: High Latitude Crabs: Biology, Management, and Economics. Alaska Sea Grant College Program. AK-SG-96-02. 713 p.
- Kuzmin SA, Akhatrin SM, Menins DT. 1999. The first findings of the snow crab *Chionoecetes opilio* (Decapoda, Majidae) in the Barents Sea. Zool Zh 77:489–491
- Kuzmin SA. 2000. Spreading of snow crab *Chionoecetes opilio* (Fabricius) in the Barents Sea. Int. Counc. Expl. Sea. CM 2000/U:21, 7 p.
- Lee JE, Chown SL. 2007 Mytilus on the move: Transport of an invasive bivalve to the Antarctic. Marine Ecology Progress Series 339:307 - 310 <https://doi.org/10.3354/meps339307>
- Lehtiniemi M, Ojaveer H, David M, Galil B, Gollasch S, McKenzie C, Minchin D, Occhipinti-Ambrogi A, Olenin S, Pederson J. 2015. Dose of truth – monitoring marine non-indigenous species to serve legislative requirements. Marine Policy 54:26-35.
- Leppäkoski E, Olenin S. 2000. Non-native species and rates of spread: lessons from the brackish Baltic Sea. Biol. Invasions, 2, 151–163.
- Leung B, Lodge DM, Finnoff D, Shogren JF, Lewis MA, Lamberti G. 2002. An ounce of prevention or a pound of cure: bioeconomic risk analysis of invasive species. Proceedings of the Royal Society B 269: 2407-2413.
- Madenjian CP, Tapanian MA, Witzel LD, Einhouse DW, Pothoven SA, Whitford HL. 2011. Evidence for predatory control of the invasive round goby. Biological Invasions 13:987–1002.
- McKenzie CH, Reid V, Lampert G, Matheson K, Minchin D, Pederson J, Brown L, Curd A, Gollasch S, Couilletquer P, Occhipinti-Ambrogi A, Simard N, Therriault TW. 2017. Alien Species Alert: *Didemnum vexillum* Kott, 2002: Invasion, impact, and control. ICES COOPERATIVE RESEARCH REPORT RAPPORT DES RECHERCHES COLLECTIVES NO. 335
- Minchin D. 2007. Rapid coastal survey for targeted alien species associated with floating pontoons in Ireland. Aquatic Invasions 2:63-70.
- Mortensen S, et al. 2021. Towards a sustainable fishery and use of cleaner fish in salmonid aquaculture.
- Mortensen S, Strand Ø. 2000. Releases and recaptures of Manila clams (*Ruditapes philippinarum*) introduced to Norway.
- Mortensen S, Strand Å, Bodvin T, Alfjorden A, Skår CK, Jelmert A, Aspán A, Sælemyr L, Naustvoll LJ, Albretsen J. 2016. Summer mortalities and detection of ostreid herpesvirus microvariant in Pacific oyster *Crassostrea gigas* in Sweden and Norway. Diseases of Aquatic Organisms 117: 171–176, <https://doi.org/10.3354/dao02944>

- Mortensen S, Bodvin T, Strand Å, Holm MW, Dolmer P. 2017. Effects of a bio-invasion of the Pacific oyster, *Crassostrea gigas* (Thunberg, 1793) in five shallow water habitats in Scandinavia. *Managing Biological Invasions* 8: 543-552.
- Mortensen S, Strand Å, Dolmer P, Laugen AT, Naustvoll L-J. 2019. Høsting av stillehavsøsters. TemaNord 2019:552, Nordisk ministerråd, ISBN 978-92-893-6425-6 (pdf) <http://dx.doi.org/10.6027/tn2019-552>, 115s. (in Norwegian).
- Mortensen S, Skår C. 2020. The surveillance and control programme for bonamiosis and marteiliosis in European flat oysters, *Ostrea edulis*, and blue mussels, *Mytilus* sp. in Norway in 2019. Rapport fra havforskningen nr 36, 2020, 16 s. <https://www.hi.no/hi/nettrapporter/rapport-fra-havforskningen-en-2020-36>
- Moseid CVS, Falkenhaug T, Slettan A. 2021. Development of a TaqMan PCR assay for the identification of the non-native copepod *Acartia tonsa*, and detection of this species in Norwegian coastal waters, *Journal of Plankton Research*, Volume 43:3, Pages 497–499, <https://doi.org/10.1093/plankt/fbab029>
- Munro ES, McIntosh RE, Weir SJ, Noguera PA, Sandilands JM, Matejusova I, Mayes AS, Smith, R. 2015. A mortality event in wrasse species (Labridae) associated with the presence of viral haemorrhagic septicaemia virus. *Journal of Fish Diseases* 38: 335–341. DOI: 10.1111/jfd.12237
- Nunkoo I, Hemmingsen W, McKenzie K, Karlsbakk E, Arneberg P. Submitted. The Barents Sea snow crab (*Chionoecetes opilio*) invasion and *Trypanosoma murmanense* infections in cod (*Gadus morhua*).
- Nunes AL, Katsanevakis A, Zenetos A, Cardoso AC. 2014. Gateways to alien invasions in the European seas. *Aquatic Invasions* 9:133-144. doi: <http://dx.doi.org/10.3391/ai.2014.9.2.02>
- Nyberg CD, Wallentinus I. 2005. Can species traits be used to predict marine macroalgal introductions? *Biological Invasions* 7: 265-279
- Næss, R. 2015. Populasjonsdynamikk og diett hos *Mnemiopsis leidyi* (Amerikansk lobemanet) UiA Bacehloroppgave i biologi. referansenummer 086-BIO
- Ojaveer H, Galil BS, Lehtiniemi M, Christoffersen M, Clink S, Florin A-B, Gruszka P, Puntilla R, Behrens J. 2015. Twenty five years of invasion: management of the round goby *Neogobius melanostomus* in the Baltic Sea. *Management of Biological Invasions* 6:329-339.
- Oliveira OMP. 2007. The presence of the ctenophore *Mnemiopsis leidyi* in the Oslofjorden and considerations on the initial invasion pathways to the North and Baltic Seas. European Research Network on Aquatic Invasive Species. *Aquatic Invasions* (2007) 2(3): 185-189.
- Orlov YI, Ivanov BG. 1978. On the introduction of Kamchatka king crab *Paralithodes camtschatica* (Decapoda: Anomura: Lithodidae) into Barents Sea. *Mar. Biol.* 48, 373-375.
- Orlov Yul, Karpevich AF. 1965. On the introduction of the commercial crab *Paralithodes camtschaticus* (Tilesius) into the Barents Sea. *Rapp. P.-V. Reun. Cons. Int. Explor. Mer* 156: 59-61.
- Ouellet P, Sainte-Marie B. 2018. Vertical distribution of snow crab (*Chionoecetes opilio*) pelagic stages in the Gulf of St. Lawrence (Canada) and effect of temperature on development and survival. *ICES J. Mar. Sci.*, 75 (2): 773-784.
- Oug E, Cohrane KJ, SundetJH, Norling K, Nilsson HC. 2011. Effects of the invasive red king crab (*Paralithodes*

- camtschaticus*) on soft-bottom fauna in Varangerfjorden, northern Norway. *Mar. Biodiv.* 41: 467-479.
- Patel S, Korsnes K, Bergh Ø, Vik-Mo F, Pedersen J, Nerland AH. 2007. Nodavirus in farmed Atlantic cod *Gadus morhua* in Norway. *Diseases of Aquatic Organisms* 77: 169–173. <https://doi.org/10.3354/dao01842>
- Pavičić M, Dragičević B, Žužul I, Vrdoljak D, Matic- Skoko S, Šegvić-Bubić T. 2020. First record of American lobster, *Homarus americanus* (H. Milne Edwards, 1837), in the Mediterranean Sea. *BioInvasions Records* 9: 83–88.
- Rech S, Borell Y, Garcia-Vazquez E. 2016. Marine litter as a vector for non-native species: What we need to know. *Marine Pollution Bulletin* 113: 40-43.
- Reise K, Olenin S, Thieltges DW. 2006. Are aliens threatening European aquatic coastal ecosystems? *Helgoland Marine Research* 60: 77–83, <https://doi.org/10.1007/s10152-006-0024-9>
- Robalo JI, Castilho R, Francisco SM, Almada F, Knutsen H, Jorde PE, Pereira AM, Almada V. 2012. Northern refugia and recent expansion in the North Sea: the case of the wrasse *Symphodus melops* (Linnaeus, 1758). *Ecology and Evolution* 2: 153–164. <https://doi.org/10.1002/ece3.77>
- Roche RC, Monnington JM, Newstead RG, Sambrook K, Griffith K, Holt RHF, Jenkins SR. 2015. Recreational vessels as a vector for marine non-natives: developing biosecurity measures and managing risk through an in-water encapsulation system. *Hydrobiologia*, 750(1): 187–199.
- Sandvik H, Dolmen D, Elven R, Falkenhaug T, Forsgren E, Hansen H, et al. 2019. Alien plants, animals, fungi and algae in Norway: an inventory of neobiota. *Biological Invasions* 21: 2997–3012. <http://dx.doi.org/10.1007/s10530-019-02058-x>
- Sandvik H, Hilmo O, Henriksen S, Elven R, Åsen PA, Hegre H, et al. 2020. Alien species in Norway: results from quantitative ecological impact assessments. *Ecological Solutions and Evidence* 1: e12006. <http://dx.doi.org/10.1002/2688-8319.12006>
- Schönherz AA, Hansen MHH, Jorgensen HBH, Berg P, Lorenzen N, Einer-Jensen K. 2012. Oral transmission as a route of infection for viral haemorrhagic septicaemia virus in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases* 35: 395-406. <https://doi.org/10.1111/j.1365-2761.2012.01358.x>
- Seljestad G, Quintela M, Faust E, Halvorsen KT, Besnier F, Jansson E, Dahle G, Knutsen H, André C, Folkvord A, Glover KA. 2020. “A cleaner-break”: Genetic divergence between geographic groups and sympatric phenotypes revealed in Ballan wrasse (*Labrus bergylta*). *Ecology and Evolution* 10: 6120–6135. <https://doi.org/10.1002/ece3.6404>
- Skaala Ø, Borstrøm R, Fiske P, Hindar K, Jensen AJ, Mo TA. 2021. Rengbueørret (*Onchorhynchus mykiss*); produksjon og interaskjoner med stedegne arter. Rapport fra Hvforskningen 2020-21.
- Slizkin AG. 1982. Distribution of snow crabs of the genus *Chionoecetes* and their habitat in the northern part of the Pacific Ocean: population dynamics and reproductive conditions of commercial invertebrates and algae in the Far Eastern Seas. *Izv Tikhookean Nauchno-Issled Inst Ryb Khoz Oceanogr*, 106:26–33 (Canadian Translation of Fisheries and Aquatic Sciences No. 5664).
- Stevens, BG. 2014. Development and biology of king crab larvae, pp. 233-259. In *King Crabs of the World. Biology and Fisheries Management*, BG Stevens (ed). CRC Press, Taylor and Francis Group. 608 p.
- Strand KO, Huserbråten M, Dagestad K-F, Mauritzen C, Grøsvik BE, Nogueira LA, Melsom A, Røhrs J. 2021.

- Potential sources of marine plastic from survey beaches in the Arctic and Northeast Atlantic. *Science of the Total Environment* 790: 1-13.
- Strand Å, Blanda E, Bodvin T, Davids JK, Jensen LF, Holm-Hansen TH, Jelmert A, Lindegarth S, Mortensen S, Moy FE, Nielsen P, Norling P, Nyberg C, Christensen HT, Vismann B, Holm MW, Hansen BW, Dolmer P. 2012. Impact of an icy winter on the Pacific oyster (*Crassostrea gigas* Thunberg, 1793) populations in Scandinavia. *Aquatic Invasions* 7: 433–440, <https://doi.org/10.3391/ai.2012.7.3.014>
- Stebbing, P, P Johnson, A Delahunty, PF Clark, T McCollin, C hale & S Clark. 2012. Reports of American lobsters, *Homarus americanus* (H. Milne Edwards, 1837), in British waters. *BioInvasions Records*, 1(1): 17-23.
- Sundal-Joys IK. 2021. Temperature tolerance and distribution of the invasive non-indigenous red algae *Agarophyton vermiculophyllum* in seagrass meadows in Norway. Master Thesis, University of Bergen. 2.11.2021.
- Sundet JH. 2014. Red king crab in the Barents Sea, pp. 485-500. In *King Crabs of the World. Biology and Fisheries Management*, BG Stevens (ed). CRC Press, Taylor and Francis Group. 608 p.
- Sundet JH, Hoel AH. 2016. The Norwegian management of an introduced species: the Arctic red king crab fishery. *Marine Policy*, 72: 278-284.
- Sundet JH, Tranang CA, Nilssen EM, Jenssen M. In prep. Stomach content of the snow crab (*Chionoecetes opilio*) from the western Barents Sea.
- Sussarellu R, Huvet A, Lapègue S, Quillen V, Lelong C, Cornette F, Jensen LF, Bierne N, Boudry P. 2015. Additive transcriptomic variation associated with reproductive traits suggest local adaptation in a recently settled population of the Pacific oyster, *Crassostrea gigas*. *BMC Genomics* 16: 808, <https://doi.org/10.1186/s12864-015-1972-8>
- Svåsand T, Grefsrud ES, Karlsen Ø, Kvamme BO, Glover K, Husa V, Kristiansen TS (Eds). 2017. Risk report Norwegian Fish Farming 2017 (in Norwegian). Institute of Marine Research. Available from: https://www.hi.no/hi/nettrapporter/risikorapport-norsk-fiskeoppdrett/2017/risikorapport_2017
- Takahashi CK, Lourenço NGGS, Lopes TF, Rall LM, Lopes CAM. 2008. BALLAST WATER: A REVIEW OF THE IMPACT ON THE WORLD PUBLIC HEALTH. *J. Venom. Anim. Toxins incl. Trop. Dis.*, 2008, 14, 3, p. 398
- Tambs-Lyche, H. 1964. *Gonionemus vertens* L. AGASSIZ (Limnomedusae) - a zoogeographical puzzle. *Sarsia* 15: 1-8
- Tiselius P, Møller LF 2017. Community cascades in a marine pelagic food web controlled by the non-visual apex predator *Mnemiopsis leidyi* J. *Plankton res.* 39: 271-279
- Tutman P, Kaparis K, Kirincic M, Pallaoro A. 2017. Floating marine litter as a raft of drifting voyages for *Planes minutus* (Crustacea: Decapoda: Grapsidae) and *Liocarcinus navigator* (Crustacea: Decapoda: Polybiidae). *Marine Pollution Bulletin* 120: 217-221.
- Verliin A, Kesler M, Svirgsden R, Taal I, Saks L, Rohtla M, Hubel, K, Eshbaum R, Vetemaa M. 2017. Invasion of round goby to the temperate salmon streams in the Baltic Sea. *Ichthyological Research* 64:155-158.
- VKM 2017. Risk assessment of fish health associated with the use of cleaner fish in aquaculture. Opinion of the Panel on Animal Health and Welfare of the Norwegian Scientific Committee for Food and Environment. VKM

report 2017:32, Oslo, Norway

(<https://vkm.no/risikovurderinger/allevurderinger/resefiskogrisikoforoverforingavsmittetiloppdrettslaks.4.d44969/>)

VKM 2019. Rueness E, Berg PR, Gulla S, Halvorsen K, Järnegren J, Malmstrøm M, Mo TA, Rimstad E, de Boer H, Eldegard K, Hindar K, Hole LR, Kausrud K, Kirkendall L, Måren I, Nilsen EB, Thorstad EB, Nielsen A, Velle G 2019. Assessment of the risk to Norwegian biodiversity from import of wrasses and other cleaner fish for use in aquaculture. Opinion of the Panel on Alien Organisms and Trade in Endangered Species (CITES) of the Norwegian Scientific Committee for Food and Environment. VKM report 2019: 15, ISBN: 978-82-8259-330-4, ISSN: 2535-4019. Oslo, Norway

Ware C, Berge J, Jelmert A, Olsen SM, Pellissier L, Wisz M, Kriticos D, Semenov G, Kwaśniewski S, Alsos IG. 2015. Biological introduction risks from shipping in a warming Arctic. *Journal of Applied Ecology*. doi: 10.1111/1365-2664.12566 [10.1111/1365-2664.12566](https://doi.org/10.1111/1365-2664.12566)

Wiegleb J, Hirsch PE, Seidel F, Rauter G, Burkhardt-Holm P. 2021. Flow, force, behaviour: assessment of a prototype hydraulic barrier for invasive fish. *Hydrobiologia*. <https://doi.org/10.1007/s10750-021-04762-z>

Williams AB. 1984. Shrimps, lobster and crabs of the Atlantic coast and eastern United States, Main to Florida, Smithsonian Institution Press, Washington DC

Woll A, Skiftesvik AB, Solevåg SE, Hansen Aas G, Bakke S, Bjelland R. 2013. Fiskestørrelse og rømming fra laksemerd. *Norsk Fiskeoppdrett August 2013* (in Norwegian).

Wonham MJ, Carlton JT, Ruiz GM, Smith LD. 2000. Fish and ships: relating dispersal frequency to success in biological invasions. *Marine Biology* 136:1111-1121.

Wrange AL, Valero J, Harketstad LS, Strand Ø, Lindegarth S, Christensen HT, Dolmer P, Kristensen PS, Mortensen S. 2010. Massive settlements of the Pacific oyster, *Crassostrea gigas*, in Scandinavia. *Biological Invasions* 12: 1145–115, <https://doi.org/10.1007/s10530-009-9535-z>

Østhagen A, Raspotnik A. 2018. Crab! How a dispute over snow crab became a diplomatic headache between Norway and the EU. *Marine policy*, 98: 58-61.

Øresland V, Ulmestrand M, Agnalt A-L, Oxby G. 2017. Recorded captures of American lobster (*Homarus americanus*) in Swedish waters and an observation of predation on the European lobster (*Homarus gammarus*). *Can. J. Fish. Aquat. Sci.*, 74: 1503-1506.

7 - Appendix 1

Table 1. Arrival of cargo vessels into Norwegian ports in the period oktober 2020- oktober 2021. * 500 arrivals =>500 arrivals. ** The annual number of arrivals is estimated for ports which recieved more than 500 arrivals per year, calculated by days/500 arrivals.

County	Port	*Arrivals	Foreign	% Foreign	Days/500 arrivals	**Annual arrivals
Agder	Flekkefjord	29	3	10 %		29
Agder	Farsund	253	44	17 %		253
Agder	Hausvik	155	50	32 %		155
Agder	Lyngdal	90	15	17 %		90
Agder	Mandal	271	29	11 %		271
Agder	Høllen	297	10	3 %		297
Agder	Flekkerøy	5	0	0 %		5
Agder	Kristiansand	500	105	21 %	189	966
Agder	Lillesand	80	15	19 %		80
Agder	Grimstad	28	11	39 %		28
Agder	Arendal	211	56	27 %		211
Agder	Eydehavn	171	82	48 %		171
Agder	Risør	10	0	0 %		10
Møre & Romsdal	Larsnes	44	0	0 %		44
Møre & Romsdal	Mjølstadneset	98	1	1 %		98
Møre & Romsdal	Fosnavåg	37	0	0 %		37
Møre & Romsdal	Ulsteinvik	99	7	7 %		99
Møre & Romsdal	Hjørungavåg	60	3	5 %		60

Møre & Romsdal	Hareid	67	1	1 %		67
Møre & Romsdal	Ørsta	138	2	1 %		138
Møre & Romsdal	Fiskarstranda	48	1	2 %		48
Møre & Romsdal	Spjelkavik	402	6	1 %		402
Møre & Romsdal	Ålesund	500	21	4 %	81	2253
Møre & Romsdal	Søvik	50	2	4 %		50
Møre & Romsdal	Steinshamn	40	0	0 %		40
Møre & Romsdal	Brattvåg	26	1	4 %		26
Møre & Romsdal	Vestnes	346	2	1 %		346
Møre & Romsdal	Molde	250	3	1 %		250
Møre & Romsdal	Elnesvågen	222	4	2 %		222
Møre & Romsdal	Harøysundet	187	3	2 %		187
Møre & Romsdal	Averøya	500	61	12 %	237	770
Møre & Romsdal	Sunnalsøra	308	37	12 %		308
Møre & Romsdal	Kristiansund	500	4	1 %	119	1534
Nordland	Brønnøysund	500	2	0 %	187	976
Nordland	Herøy	500	0	0 %	346	527
Nordland	Sandnessjøen	490	0	0 %		490
Nordland	Mosjøen	182	71	39 %		182
Nordland	Mo i Rana	500	156	31 %	238	767
Nordland	Halsa Meløy	500	21	4 %	169	1080
Nordland	Glomfjord	319	104	33 %		319
Nordland	Ørnes	312	0	0 %		312

Nordland	Fauske	165	0	0 %		165
Nordland	Rognan	35	4	11 %		35
Nordland	Bodø	500	0	0 %	144	1267
Nordland	Kjøpsvik	169	28	17 %		169
Nordland	Hekkelstrand	500	1	0 %	286	638
Nordland	Narvik	376	196	52 %		376
Nordland	Lødingen	280	2	1 %		280
Nordland	Svolvær	371	0	0 %		371
Nordland	Leknes	159	1	1 %		159
Nordland	Stokkmarknes	500	8	2 %	281	649
Nordland	Sortland	500	7	1 %	128	1426
Nordland	Myre	486	5	1 %		486
Nordland	Andenes	50	0	0 %		50
Oslo	Oslo	500	223	45 %	88	2074
Rogaland	Ølen	167	5	3 %		167
Rogaland	Haugesund	500	48	10 %	32	5703
Rogaland	Storasund	500	9	2 %	153	1193
Rogaland	Husøya	500	78	16 %	100	1825
Rogaland	Skudeneshavn	105	17	16 %		105
Rogaland	Kårstø	11	0	0 %		11
Rogaland	Sauda	291	94	32 %		291
Rogaland	Forsand	332		0 %		332
Rogaland	Sandnes	408	71	17 %		408

Rogaland	Stavanger anch	139	42	30 %		139
Rogaland	Mekjarvik	129	18	14 %		129
Rogaland	Tananger	500	180	36 %	124	1472
Rogaland	Sirevåg	48	8	17 %		48
Rogaland	Egersund	481	142	30 %		481
Rogaland	Jøssingfjord	218	105	48 %		218
Rogaland	Breiviken	500	109	22 %	148	1233
Rogaland	Stavanger	500	89	18 %	225	811
Rogaland	Dusavik	216	14	6 %		216
Svalbard	Ny Ålesund	20	0	0 %		20
Svalbard	Longyearbyen	44	0	0 %		44
Troms & Finnmark	Bergneset	370	11	3 %		370
Troms & Finnmark	Harstad	500	2	0 %	193	946
Troms & Finnmark	Finnsnes	320	0	0 %		320
Troms & Finnmark	Tromsø	500	4	1 %	67	2724
Troms & Finnmark	Øksfjord	350	0	0 %		350
Troms & Finnmark	Alta	127	0	0 %		127
Troms & Finnmark	Rypefjord	500	1	0 %	196	931
Troms & Finnmark	Hammerfest	500	0	0 %	165	1106
Troms & Finnmark	Melkeøya	9	0	0 %		9
Troms & Finnmark	Havøysund	396	0	0 %		396
Troms & Finnmark	Honningsvåg	197	0	0 %		197
Troms & Finnmark	Mehamn	42	0	0 %		42

Troms & Finnmark	Båtsfjord	204	45	22 %		204
Troms & Finnmark	Vardø	20	1	5 %		20
Troms & Finnmark	Vadsø	19	0	0 %		19
Troms & Finnmark	Kirkenes	121	41	34 %		121
Trøndelag	Nordskaget	19	0	0 %		19
Trøndelag	Orkanger	298	71	24 %		298
Trøndelag	Trondheim	500	21	4 %	217	841
Trøndelag	Verdal	238	56	24 %		238
Trøndelag	Lysøysund	45	0	0 %		45
Trøndelag	Kjerkeholmen	94	0	0 %		94
Trøndelag	Sandviksberget	500	0	0 %	323	565
Trøndelag	Namsos	224	0	0 %		224
Trøndelag	Rørvik	500	0	0 %	169	1080
Vestfold og Telemark	Kragerø Anch	33	16	48 %		33
Vestfold og Telemark	Kragerø	446	153	34 %		446
Vestfold og Telemark	Langesund	141	0	0 %		141
Vestfold og Telemark	Brevik	500	183	37 %	265	689
Vestfold og Telemark	Rafnes	131	100	76 %		131
Vestfold og Telemark	Porsgrunn	500	292	58 %	194	941

Vestfold og Telemark	Larvik	395	90	23 %		395
Vestfold og Telemark	Sandefjord	4	0	0 %		4
Vestfold og Telemark	Slagen	0	0	0 %		0
Vestfold og Telemark	Horten	196	125	64 %		196
Vestfold og Telemark	Holmestrand	34	30	88 %		34
Vestland	Hanøytangen	86	8	9 %		86
Vestland	Ågotnes	458	45	10 %		458
Vestland	Skipavik	216	49	23 %		216
Vestland	Mongstad	410	26	6 %		410
Vestland	Sture	2	0	0 %		2
Vestland	Storabø	156	5	3 %		156
Vestland	Torangsvåg	415	3	1 %		415
Vestland	Rubbestadneset	64	1	2 %		64
Vestland	Stord havn	152	26	17 %		152
Vestland	Odda	249	152	61 %		249
Vestland	Husnes	173	15	9 %		173
Vestland	Høylandsbygd	15	0	0 %		15
Vestland	Sløvåg anch	68	16	24 %		68
Vestland	Sløvåg	427	87	20 %		427
Vestland	Leirvik Hyllestad	103	2	2 %		103

Vestland	Høyanger	446	26	6 %		446
Vestland	Årdalstangen	197	42	21 %		197
Vestland	Florø	500	39	8 %	73	2500
Vestland	Svelgen	209	42	20 %		209
Vestland	Kalvåg	96	0	0 %		96
Vestland	Måløy	500	17	3 %	70	2607
Vestland	Raudeberg	93	2	2 %		93
Vestland	Kolsnes	39	3	8 %		39
Vestland	Eikefet	500	64	13 %		500
Vestland	Bergen	500	12	2 %	94	1941
Vestland	Leirvik	265	1	0 %		265
Viken	Drammen	473	250	53 %		473
Viken	Drøbak	1	0	0 %		1
Viken	Kambo	97	46	47 %		97
Viken	Moss	306	100	33 %		306
Viken	Fredrikstad	500	235	47 %	165	1106
Viken	Hvaler	0	0	0 %		0
Viken	Halden	112	18	16 %		112

Table 2. Arrival of tugs & special crafts into Norwegian ports in the period oktober 2020- oktober 2021. * 500 arrivals =>500 arrivals. ** The annual number of arrivals is estimated for ports which recieved more than 500 arrivals per year, calculated by days/500 arrivals.

County	Port	*Arrivals	Foreign	% Foreign	Days/500 arrivals	*Annual arrivals
Agder	Flekkefjord	135	3	2 %		135
Agder	Farsund	500	5	1 %	158	1155
Agder	Hausvik	24	0	0 %		24
Agder	Lyngdal	51	3	6 %		51
Agder	Mandal	374	6	2 %		374
Agder	Høllen	218	0	0 %		218
Agder	Flekkerøy	261	0	0 %		261
Agder	Kristiansand	500	3	1 %	67	2724
Agder	Lillesand	385	0	0 %		385
Agder	Grimstad	189	1	1 %		189
Agder	Arendal	500	2	0 %	167	1093

Agder	Eydehavn	140	3	2 %		140
Agder	Risør	116	3	3 %		116
Møre & Romsdal	Larsnes	500	0	0 %	252	724
Møre & Romsdal	Mjølstadneset	99	2	2 %		99
Møre & Romsdal	Fosnavåg	254	1	0 %		254
Møre & Romsdal	Ulsteinvik	191	7	4 %		191
Møre & Romsdal	Hjørungavåg	22	0	0 %		22
Møre & Romsdal	Hareid	114	4	4 %		114
Møre & Romsdal	Ørsta	26	2	8 %		26
Møre & Romsdal	Fiskarstranda	86	2	2 %		86
Møre & Romsdal	Spjelkavik	124	5	4 %		124
Møre & Romsdal	Ålesund	500	2	0 %	48	3802
Møre & Romsdal	Søvik	114	0	0 %		114
Møre & Romsdal	Steinshamn	241	0	0 %		241
Møre & Romsdal	Brattvåg	269	0	0 %		269
Møre & Romsdal	Vestnes	500	3	1 %		500
Møre & Romsdal	Molde	500	1	0 %	326	560
Møre & Romsdal	Elnesvågen	87	0	0 %		87
Møre & Romsdal	Harøysundet	96	0	0 %		96
Møre & Romsdal	Averøya	263	2	1 %		263
Møre & Romsdal	Sunnalsøra	86	0	0 %		86
Møre & Romsdal	Kristiansund	500	0	0 %	33	5530
Nordland	Brønnøysund	500	2	0 %	213	857

Nordland	Herøy	500	0	0 %	194	941
Nordland	Sandnessjøen	500	0	0 %	53	3443
Nordland	Mosjøen	71	0	0 %		71
Nordland	Mo i Rana	111	2	2 %		111
Nordland	Halsa Meløy	87	0	0 %		87
Nordland	Glomfjord	61	0	0 %		61
Nordland	Ørnes	74	0	0 %		74
Nordland	Fauske	110	0	0 %		110
Nordland	Rognan	19	1	5 %		19
Nordland	Bodø	500	3	1 %	115	1587
Nordland	Kjøpsvik	12	0	0 %		12
Nordland	Hekkelstrand	335	0	0 %		335
Nordland	Narvik	500	0	0 %	288	634
Nordland	Lødingen	500	0	0 %	147	1241
Nordland	Svolvær	500	0	0 %	293	623
Nordland	Leknes	68	0	0 %		68
Nordland	Stokkmarknes	466	0	0 %		466
Nordland	Sortland	500	0	0 %	93	1962
Nordland	Myre	500	0	0 %	256	713
Nordland	Andenes	299	0	0 %		299
Oslo	Oslo	500	33	7 %	8	22813
Rogaland	Ølen	155	2	1 %		155
Rogaland	Haugesund	500	17	3 %	66	2765

Rogaland	Storasund	500	1	0 %	140	1304
Rogaland	Husøya	500	16	3 %	261	699
Rogaland	Skudeneshavn	139	0	0 %		139
Rogaland	Kårstø	500	0	0 %	116	1573
Rogaland	Sauda	189	0	0 %		189
Rogaland	Forsand	37	0	0 %		37
Rogaland	Sandnes	130	1	1 %		130
Rogaland	Stavanger anch	500	26	5 %	309	591
Rogaland	Mekjarvik	484	27	6 %		484
Rogaland	Tananger	500	14	3 %	47	3883
Rogaland	Sirevåg	31	0	0 %		31
Rogaland	Egersund	500	3	1 %	306	596
Rogaland	Jøssingfjord	89	1	1 %		89
Rogaland	Breiviken	289	0	0 %		289
Rogaland	Stavanger	500	16	3 %	57	3202
Rogaland	Dusavik	500	11	2 %	105	1738
Svalbard	Ny Ålesund	153	0	0 %		153
Svalbard	Longyearbyen	500	0	0 %	163	1120
Troms & Finnmark	Bergneset	14	0	0 %		14
Troms & Finnmark	Harstad	500	1	0 %	65	2808
Troms & Finnmark	Finnsnes	447	0	0 %		447
Troms & Finnmark	Tromsø	500	0	0 %	38	4803
Troms & Finnmark	Øksfjord	74	0	0 %		74

Troms & Finnmark	Alta	348	0	0 %		348
Troms & Finnmark	Rypefjord	500	0	0 %	143	1276
Troms & Finnmark	Hammerfest	500	1	0 %	101	1807
Troms & Finnmark	Melkeøya	122	0	0 %		122
Troms & Finnmark	Havøysund	431	0	0 %		431
Troms & Finnmark	Honningsvåg	500	0	0 %	282	647
Troms & Finnmark	Mehamn	188	0	0 %		188
Troms & Finnmark	Båtsfjord	277	0	0 %		277
Troms & Finnmark	Vardø	500	0	0 %	236	773
Troms & Finnmark	Vadsø	27	5	19 %		27
Troms & Finnmark	Kirkenes	500	13	3 %	278	656
Trøndelag	Nordskaget	39	0	0 %		39
Trøndelag	Orkanger	45	2	4 %		45
Trøndelag	Trondheim	500	1	0 %	123	1484
Trøndelag	Verdal	156	12	8 %		156
Trøndelag	Lysøysund	321	1	0 %		321
Trøndelag	Kjerkeholmen	496	0	0 %		496
Trøndelag	Sandviksberget	330	0	0 %		330
Trøndelag	Namsos	148	0	0 %		148
Trøndelag	Rørvik	500	1	0 %	107	1706
Vestfold og Telemark	Kragerø	500	4	1 %	128	1426
Vestfold og Telemark	Langesund	500	0	0 %	54	3380

Vestfold og Telemark	Brevik	500	4	1 %	13	14038
Vestfold og Telemark	Rafnes	500	0	0 %	139	1313
Vestfold og Telemark	Porsgrunn	500	1	0 %	90	2028
Vestfold og Telemark	Larvik	124	2	2 %		124
Vestfold og Telemark	Sandefjord	139	1	1 %		139
Vestfold og Telemark	Slagen	500	18	4 %	289	631
Vestfold og Telemark	Horten	500	1	0 %	176	1037
Vestfold og Telemark	Holmestrand	153	0	0 %		153
Vestland	Hanøytangen	271	6	2 %		271
Vestland	Ågotnes	500	12	2 %	270	676
Vestland	Skipavik	411	6	1 %		411
Vestland	Mongstad	500	6	1 %	47	3883
Vestland	Sture	500	0	0 %	210	869
Vestland	Storabø	132	2	2 %		132
Vestland	Torangsvåg	417	1	0 %		417
Vestland	Rubbestadneset	269	2	1 %		269
Vestland	Stord havn	340	6	2 %		340
Vestland	Odda	43	0	0 %		43
Vestland	Husnes	111	0	0 %		111

Vestland	Høylandsbygd	292	3	1 %		292
Vestland	Sløvåg anch	288	4	1 %		288
Vestland	Sløvåg	500	2	0 %	133	1372
Vestland	Leirvik Hyllestad	284	0	0 %		284
Vestland	Høyanger	105	2	2 %		105
Vestland	Årdalstangen	66	0	0 %		66
Vestland	Florø	500	7	1 %	46	3967
Vestland	Svelgen	51	0	0 %		51
Vestland	Kalvåg	182	0	0 %		182
Vestland	Måløy	500	9	2 %	131	1393
Vestland	Raudeberg	269	2	1 %		269
Vestland	Kolsnes	500	0	0 %	90	2028
Vestland	Eikefet	40	0	0 %		40
Vestland	Bergen	500	23	5 %	54	3380
Vestland	Leirvik	500	3	1 %	310	589
Viken	Drammen	310	6	2 %		310
Viken	Drøbak	387	0	0 %		387
Viken	Kambo	205	3	1 %		205
Viken	Moss	500	0	0 %	283	645
Viken	Fredrikstad	500	3	1 %	195	936
Viken	Hvaler	500	5	1 %	183	997
Viken	Halden	202	5	2 %		202

Table 3. Arrival of tankers into Norwegian ports in the period oktober 2020- oktober 2021. * 500 arrivals =>500 arrivals. ** The annual number of arrivals is estimated for ports which recieved more than 500 arrivals per year, calculated by days/500 arrivals.

County	Port	*Arrivals	Foreign	% Foreign	Days/500 arrivals	**Annual arrivals
Agder	Flekkefjord	1	0	0 %		1
Agder	Farsund	32	8	25 %		32
Agder	Hausvik	2	0	0 %		2
Agder	Lyngdal	1	0	0 %		1
Agder	Mandal	4	0	0 %		4
Agder	Høllen	1	0	0 %		1
Agder	Flekkerøy	0	0	0 %		0
Agder	Kristiansand	134	0	0 %		134
Agder	Lillesand	0	0	0 %		0

Agder	Grimstad	0	0	0 %		0
Agder	Arendal	3	0	0 %		3
Agder	Eydehavn	0	0	0 %		0
Agder	Risør	2	0	0 %		2
Møre & Romsdal	Larsnes	2	0	0 %		2
Møre & Romsdal	Mjølstadneset	5	0	0 %		5
Møre & Romsdal	Fosnavåg	18	0	0 %		18
Møre & Romsdal	Ulsteinvik	11	0	0 %		11
Møre & Romsdal	Hjørungavåg	4	0	0 %		4
Møre & Romsdal	Hareid	5	0	0 %		5
Møre & Romsdal	Ørsta	1	0	0 %		1
Møre & Romsdal	Fiskarstranda	14	0	0 %		14
Møre & Romsdal	Spjelkavik	28	15	54 %		28
Møre & Romsdal	Ålesund	500	0	0 %	319	572
Møre & Romsdal	Søvik	10	0	0 %		10
Møre & Romsdal	Steinshamn	6	0	0 %		6
Møre & Romsdal	Brattvåg	6	0	0 %		6
Møre & Romsdal	Vestnes	15	3	20 %		15
Møre & Romsdal	Molde	4	0	0 %		4
Møre & Romsdal	Elnesvågen	216	150	69 %		216
Møre & Romsdal	Harøysundet	5	0	0 %		5
Møre & Romsdal	Averøya	123	13	11 %		123
Møre & Romsdal	Sunnalsøra	34	0	0 %		34

Møre & Romsdal	Kristiansund	496	16	3 %		496
Nordland	Brønnøysund	170	0	0 %		170
Nordland	Herøy	55	0	0 %		55
Nordland	Sandnessjøen	97	3	3 %		97
Nordland	Mosjøen	73	22	30 %		73
Nordland	Mo i Rana	13	0	0 %		13
Nordland	Halsa Meløy	67	2	3 %		67
Nordland	Glomfjord	26	12	46 %		26
Nordland	Ørnes	3	0	0 %		3
Nordland	Fauske	0	0	0 %		0
Nordland	Rognan	1	0	0 %		1
Nordland	Bodø	144	11	8 %		144
Nordland	Kjøpsvik	2	0	0 %		2
Nordland	Hekkelstrand	1	0	0 %		1
Nordland	Narvik	7	0	0 %		7
Nordland	Lødingen	11	0	0 %		11
Nordland	Svolvær	9	0	0 %		9
Nordland	Leknes	1	0	0 %		1
Nordland	Stokkmarknes	73	1	1 %		73
Nordland	Sortland	3	0	0 %		3
Nordland	Myre	80	0	0 %		80
Nordland	Andenes	9	1	11 %		9
Oslo	Oslo	191	82	43 %		191

Rogaland	Ølen	14	0	0 %		14
Rogaland	Haugesund	500	44	9 %	226	808
Rogaland	Storasund	150	0	0 %		150
Rogaland	Husøya	98	42	43 %		98
Rogaland	Skudeneshavn	2	0	0 %		2
Rogaland	Kårstø	500	457	91 %		500
Rogaland	Sauda	4	0	0 %		4
Rogaland	Forsand	0	0	0 %		0
Rogaland	Sandnes	0	0	0 %		0
Rogaland	Stavanger anch	36	14	39 %		36
Rogaland	Mekjarvik	11	1	9 %		11
Rogaland	Tananger	241	73	30 %		241
Rogaland	Sirevåg	1	0	0 %		1
Rogaland	Egersund	48	9	19 %		48
Rogaland	Jøssingfjord	3	0	0 %		3
Rogaland	Breviken	0	0	0 %		0
Rogaland	Stavanger	159	60	38 %		159
Rogaland	Dusavik	83	20	24 %		83
Svalbard	Ny Ålesund	3	0	0 %		3
Svalbard	Longyearbyen	18	0	0 %		18
Troms & Finnmark	Bergneset	66	0	0 %		66
Troms & Finnmark	Harstad	193	7	4 %		193
Troms & Finnmark	Finnsnes	25	0	0 %		25

Troms & Finnmark	Tromsø	500	9	2 %	293	623
Troms & Finnmark	Øksfjord	2	0	0 %		2
Troms & Finnmark	Alta	35	4	11 %		35
Troms & Finnmark	Rypefjord	65	6	9 %		65
Troms & Finnmark	Hammerfest	58	0	0 %		58
Troms & Finnmark	Melkeøya	5	2	40 %		5
Troms & Finnmark	Havøysund	0	0	0 %		0
Troms & Finnmark	Honningsvåg	11	0	0 %		11
Troms & Finnmark	Mehamn	7	0	0 %		7
Troms & Finnmark	Båtsfjord	26	0	0 %		26
Troms & Finnmark	Vardø	1	0	0 %		1
Troms & Finnmark	Vadsø	0	0	0 %		0
Troms & Finnmark	Kirkenes	29	1	3 %		29
Trøndelag	Nordskaget	48	2	4 %		48
Trøndelag	Orkanger	3	0	0 %		3
Trøndelag	Trondheim	170	41	24 %		170
Trøndelag	Verdal	14	1	7 %		14
Trøndelag	Lysøysund	155	1	1 %		155
Trøndelag	Kjerkeholmen	15	0	0 %		15
Trøndelag	Sandviksberget	7	0	0 %		7
Trøndelag	Namsos	0	0	0 %		0
Trøndelag	Rørvik	119	0	0 %		119

Vestfold og Telemark	Kragerø	1	0	0 %		1
Vestfold og Telemark	Langesund	1	0	0 %		1
Vestfold og Telemark	Brevik	2	1	50 %		2
Vestfold og Telemark	Rafnes	500	402	80 %	337	542
Vestfold og Telemark	Porsgrunn	233	180	77 %		233
Vestfold og Telemark	Larvik	25	24	96 %		25
Vestfold og Telemark	Sandefjord	91	79	87 %		91
Vestfold og Telemark	Slagen	452	196	43 %		452
Vestfold og Telemark	Horten	1	1	100 %		1
Vestfold og Telemark	Holmestrand	0	0	0 %		0
Vestland	Hanøytangen	9	0	0 %		9
Vestland	Ågotnes	101	1	1 %		101
Vestland	Skipavik	17	0	0 %		17
Vestland	Mongstad	500	316	63 %	70	2607
Vestland	Sture	203	183	90 %		203
Vestland	Storabø	35	0	0 %		35
Vestland	Torangsvåg	75	0	0 %		75
Vestland	Rubbestadneset	2	1	50 %		2

Vestland	Stord havn	7	0	0 %		7
Vestland	Odda	21	11	52 %		21
Vestland	Husnes	3	0	0 %		3
Vestland	Høylandsbygd	0	0	0 %		0
Vestland	Sløvåg anch	5	1	20 %		5
Vestland	Sløvåg	37	6	16 %		37
Vestland	Leirvik Hyllestad	1	0	0 %		1
Vestland	Høyanger	28	0	0 %		28
Vestland	Årdalstangen	0	0	0 %		0
Vestland	Florø	286	52	18 %		286
Vestland	Svelgen	29	6	21 %		29
Vestland	Kalvåg	2	0	0 %		2
Vestland	Måløy	405	4	1 %		405
Vestland	Raudeberg	1	0	0 %		1
Vestland	Kolsnes	164	0	0 %		164
Vestland	Eikefet	7	2	29 %		7
Vestland	Bergen	473	9	2 %		473
Vestland	Leirvik	8	0	0 %		8
Viken	Drammen	53	50	94 %		53
Viken	Drøbak	0	0	0 %		0
Viken	Kambo	16	1	6 %		16
Viken	Moss	3	2	67 %		3

Viken	Fredrikstad	473	206	44 %		473
Viken	Hvaler	0	0	0 %		0
Viken	Halden	4	2	50 %		4



HAVFORSKNINGSINSTITUTTET

Postboks 1870 Nordnes

5817 Bergen

Tlf: 55 23 85 00

E-post: post@hi.no

www.hi.no